

Dynamics of Classical and Quantum Information on Spin-chains with Multiple Interactions

Ahmad Redwan¹, Abdel-Haleem Abdel-Aty^{2,3,*}, Nour Zidan^{4,5} and Tarek El-Shahat¹

¹ Mathematics Department, Faculty of Science, Al-Azhar University, 71524 Assiut, Egypt

² Physics Department, College of Sciences, University of Bisha, Bisha, Saudi Arabia

³ Physics Department, Faculty of Science, Al-Azhar University, 71524 Assiut, Egypt

⁴ Mathematics Department, Faculty of Science, Sohag University, 82424 Soahg, Egypt

⁵ Department of Mathematics, College of Science, Jouf University, Sakaka, Saudi Arabia

Received: 1 Jan. 2018, Revised: 28 Mar. 2018, Accepted: 2 Apr. 2018

Published online: 1 May 2018

Abstract: The classical and quantum information is very sensitive to any internal and external noise. In this paper we investigated the effect of the Calogero-Moser (CM) type interaction, Dzyaloshinskii-Moriya (DM) interaction, thermalization and uniform and nonuniform magnetic fields on the dynamics of both classical and quantum information. The spin-chain model is used to describe the system in the thermal state. The generated correlation is quantified by the Shannon entropy for the classical information and the Von Neuman entropy for the quantum one. The obtained results showed that the quantum information is more influenced by the distance between the spins.

Keywords: Shannon entropy, Von Neuman entropy, Spin-chain systems, thermal state, uniform and nonuniform magnetic field.

1 Introduction

The recent years witnessed a big revolution in the area of quantum information. The quantum computer may will be available commercially shortly [1,2], also the quantum communication and quantum networks have very speed progress and also will be turned on very soon [3,4]. Classical and quantum information are very sensitive to the internal and external noises [5,6,7,8,9,10].

The spin-chain model is used to simulate the quantum system perfectly from the quantum information, computation, communication [11], meteorology [12] and quantum thermodynamics [13]. Classical and quantum information in XX spins [14], the thermal state in XY spin-chain model is studied in [15] and in the time domain is studied by Burrell & Osborne [16]. Sarandy studied the dynamics of the classical information and the quantum discord in XXZ spin-chains [17]. Also, the classical and quantum correlations dynamics over the XXX [18] and XYZ [19] spin-chain models are investigated. The effect of the internal (spin-orbit coupling and Dzyaloshinskii-Moriya (DM) interaction) and external (uniform and non-uniform magnetic fields) interactions

on the classical and quantum information of spin-chains is studied by several groups [20,21,22].

The previous studies showed that the long-range interactions play an important role in the area of quantum information [23,24,25]. The quantum information is quantified using quantum discord between to spin with the effect of the Calogero-Moser type interaction [25]. Moreover, the possibility of using the long range interactions between the spins as quantum communication channels is investigated [26,27,28,29]. The entanglement under the effect of long-range interactions is investigated in several papers. Sultani et. al., studied the quantum entanglement between two spins in the Ising model with an added Dzyaloshinsky-Moriya (DM) interaction and in the presence of the transverse magnetic field [30,31]. The quantum state transfer and entanglement renormalization using Long-Range interactions is investigated by Eldredge et. al. [32]. Our motivation in this paper is to extend the previous work by studying the effect of long range interaction with different types of noise in the dynamics of the classical and quantum information of the spin chain system.

* Corresponding author e-mail: halimaty@gmail.com, amabdelaty@ub.edu.sa

This paper is organized as follow: the mathematical model and it is solution is addressed in section (2). The results and discussion is presented in section (2). We conclude our results in section (4).

2 The Model and it's solution for a two-qubit spin-chain

The Hamiltonian H for a two-qubit anisotropic Heisenberg model with z-component interaction parameter D_z is

$$H = J_z \sigma_1^z \sigma_2^z + D_z (\sigma_1^x \sigma_2^y - \sigma_1^y \sigma_2^x) + (B + b) \sigma_1^z + (B - b) \sigma_2^z + J(R) (\sigma_1^x \sigma_2^x + \sigma_1^y \sigma_2^y) \quad (1)$$

where J_z is the real coupling constant, D_z is the z-component DM interaction parameter, and $\sigma^i (i = x, y, z)$ are Pauli matrices. B is the homogeneous part of the magnetic field and b show the inhomogeneous. The DM-interaction and external magnetic fields are thought to be along the z-direction. $J(R)$ is the spin interaction coupling which will be characterized as far as CM interactions. All the parameters are dimensionless. In matrix form we can write the system Hamiltonian Eq.1 as follows:

$$H = \begin{pmatrix} 2B + J_z & 0 & 0 & 0 \\ 0 & 2b - J_z & 2(-iD_z + J(R)) & 0 \\ 0 & 2(iD_z + J(R)) & -2b - J_z & 0 \\ 0 & 0 & 0 & J_z - 2B \end{pmatrix} \quad (2)$$

We consider the first type of the CM model, which is a version of the Haldane-Shastry model with exchange interaction $J = 1/R^2$.

The eigenvalues for this type of CM interaction are given by:

$$\begin{aligned} E_1 &= 2B + J_z \\ E_2 &= J_z - 2B \\ E_3 &= 2\sqrt{b^2 + Dz^2 + G^2} - J_z \\ E_4 &= -2\sqrt{b^2 + Dz^2 + G^2} - J_z \\ |\psi_1\rangle &= |00\rangle \\ |\psi_2\rangle &= |11\rangle \\ |\psi_3\rangle &= \frac{1}{\sqrt{1 + MM^*}} (M|01\rangle + |10\rangle) \\ |\psi_4\rangle &= \frac{1}{\sqrt{1 + NN^*}} (M|01\rangle + |10\rangle) \end{aligned}$$

where

$$\begin{aligned} M &= \frac{\sqrt{b^2 + Dz^2 + G^2} + b}{G + iDz} \\ N &= \frac{b - \sqrt{b^2 + Dz^2 + G^2}}{G + iDz} \\ G &= \frac{1}{R^2} \end{aligned} \quad (3)$$

Once we get the eigenvalues and eigenvectors of the system Hamiltonian, we can obtain the dynamics of the density operator in the thermal state as follows:

$$\rho(T) = \frac{1}{z} \sum_i (\exp(-\frac{E_i}{T})) |\psi_i\rangle \langle \psi_i| \quad (4)$$

where $z = \text{Tr}(\rho(T))$. In the basis $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ we can obtain the density operator of the system as follows:

$$\rho(T) = \frac{1}{z} \begin{pmatrix} \rho_{11} & 0 & 0 & 0 \\ 0 & \rho_{22} & \rho_{23} & 0 \\ 0 & \rho_{32} & \rho_{33} & 0 \\ 0 & 0 & 0 & \rho_{44} \end{pmatrix} \quad (5)$$

has matrix elements given in Eq. 5 as follows:

$$\begin{aligned} \rho_{11} &= e^{2B + J_z} \\ \rho_{22} &= e^{-J_z} \left(\frac{b \sinh(2\sqrt{b^2 + Dz^2 + G^2})}{\sqrt{b^2 + Dz^2 + G^2}} + \cosh(2\sqrt{b^2 + Dz^2 + G^2}) \right) \\ \rho_{23} &= \frac{e^{-J_z} (G - iDz) \sinh(2\sqrt{b^2 + Dz^2 + G^2})}{\sqrt{b^2 + Dz^2 + G^2}} \\ \rho_{32} &= \frac{e^{-J_z} (G + iDz) \sinh(2\sqrt{b^2 + Dz^2 + G^2})}{\sqrt{b^2 + Dz^2 + G^2}} \\ \rho_{33} &= e^{-J_z} \left(\cosh(2\sqrt{b^2 + Dz^2 + G^2}) - \frac{b \sinh(2\sqrt{b^2 + Dz^2 + G^2})}{\sqrt{b^2 + Dz^2 + G^2}} \right) \\ \rho_{44} &= e^{J_z - 2B} \\ z &= 2e^{-J_z} \cosh(2\sqrt{b^2 + Dz^2 + G^2}) + 2e^{J_z} \cosh(2B) \end{aligned} \quad (6)$$

By using the $\rho(T)$ as in eq.5 it is easy to study the dynamics of the classical and quantum information using the Shannon and Von Neumann entropies as detailed in the following section.

3 Results and Discussion

In this section we studied the dynamics of the classical (Shannon entropy) and quantum information (Von Neumann entropy) over the system and it is dependence on the isotropic and anisotropic interactions.

In fig. (1) we investigated the dynamics of the Shannon and Von Neumann entropies as a function of the distance between the qubits "R" and the temperature "T" in the presence of several interactions (Spin-orbit interactions, uniform and non-uniform magnetic fields, Calogero-Moser (CM) type interaction, Dzyaloshinskii-Moriya (DM) interaction) where $J_z = D_z = B = b = 1$ where (a) for the Shannon entropy and (b) for the Von Neumann entropy. We can see from fig. (1)-a that the classical information is started from $S_H = 0.5$ at $R = 0$ and continued until $R = 0.2$, the Shannon entropy increased suddenly and reached its maximum value $S_H = 1$ at $R = 0.4$. Also, we can see that by increasing the temperature the disturbance of the system increased and consequently the classical information increased. In fig. (1)-b the quantum

information has the same behavior with the classical information, except the initial period of the figure, where the quantum information is initially started equal to zero and with distance increased the Von Neumann entropy increased and reached the maximum value as in figure (1)-a. Also, the temperature have the same effect as with classical information.

4 Conclusion

In this paper a model of spin-chain model with several types of interactions is designed and analytically solved. The dynamics of the classical and quantum correlation are investigated under the effect of the system parameters. It is found that the long range interactions have bad effect on the both quantum and classical information but by increasing the temperature the classical and quantum information recovered again.

References

- [1] Kings et al. Computational power consumption and speedup. Technical report, D-Wave Technical Report 14-1003A-C, 2017.
- [2] V. S. Denchev, S. Boixo, S. V. Isakov, N. Ding, R. Babbush, V. Smelyanskiy, J. Martinis, and H. Neven. What is the computational value of finite-range tunneling? *Phys. Rev. X*, **6**:031015, Aug (2016).
- [3] Juan Yin, Yuan Cao, Yu-Huai Li, Sheng-Kai Liao, Liang Zhang, Ji-Gang Ren, Wen-Qi Cai, Wei-Yue Liu, Bo Li, Hui Dai, Guang-Bing Li, Qi-Ming Lu, Yun-Hong Gong, Yu Xu, Shuang-Lin Li, Feng-Zhi Li, Ya-Yun Yin, Zi-Qing Jiang, Ming Li, Jian-Jun Jia, Ge Ren, Dong He, Yi-Lin Zhou, Xiao-Xiang Zhang, Na Wang, Xiang Chang, Zhen-Cai Zhu, Nai-Le Liu, Yu-Ao Chen, Chao-Yang Lu, Rong Shu, Cheng-Zhi Peng, Jian-Yu Wang, and Jian-Wei Pan. Satellite-based entanglement distribution over 1200 kilometers. *Science*, **356**:1140–1144, (2017).
- [4] Juan Yin, Yuan Cao, Yu-Huai Li, Ji-Gang Ren, Sheng-Kai Liao, Liang Zhang, Wen-Qi Cai, Wei-Yue Liu, Bo Li, Hui Dai, Ming Li, Yong-Mei Huang, Lei Deng, Li Li, Qiang Zhang, Nai-Le Liu, Yu-Ao Chen, Chao-Yang Lu, Rong Shu, Cheng-Zhi Peng, Jian-Yu Wang, and Jian-Wei Pan. Satellite-to-ground entanglement-based quantum key distribution. *Phys. Rev. Lett.*, **119**:200501, (2017).
- [5] Dan Marinescu. *Classical and Quantum Information: 1st Edition*. Elsevier, Holland, (2011).
- [6] T. Yu and J.H. Eberly. Sudden death of entanglement: Classical noise effects. *Optics Communications*, **264**:393–397, (2006).
- [7] I. Devetak and P. W. Shor. The capacity of a quantum channel for simultaneous transmission of classical and quantum information. *Communications in Mathematical Physics*, **256**:287–303, (2005).
- [8] B. Schumacher and M. D. Westmoreland. Sending classical information via noisy quantum channels. *Phys. Rev. A*, **56**:131–138, (1997).
- [9] F. Caruso, S. F. Huelga, and M. B. Plenio. Noise-enhanced classical and quantum capacities in communication networks. *Phys. Rev. Lett.*, **105**:190501, (2010).
- [10] Ch. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal. Entanglement-assisted classical capacity of noisy quantum channels. *Phys. Rev. Lett.*, **83**:3081–3084, Oct (1999).
- [11] A. Abdel-Aty, L. Y. Cheong, N. Zakaria, and N. Metwally. Quantum network via partial entangled state. *Journal of Communications*, **9**:379–384, (2014).
- [12] Michael Skotiniotis, Pavel Sekatski, and Wolfgang Dr. Quantum metrology for the ising hamiltonian with transverse magnetic field. *New Journal of Physics*, **17**:073032, (2015).
- [13] Th. P. Le, J. Levinsen, K. Modi, M. M. Parish, and F. A. Pollock. Spin-chain model of a many-body quantum battery. *Phys. Rev. A*, **97**:022106, (2018).
- [14] L. Henderson and V. Vedral. Classical, quantum and total correlations. *Journal of Physics A: Mathematical and General*, **34**:6899, (2001).
- [15] J. Maziero, H. C. Guzman, L. C. Céleri, M. S. Sarandy, and R. M. Serra. Quantum and classical thermal correlations in the XY spin- $\frac{1}{2}$ chain. *Phys. Rev. A*, **82**:012106, (2010).
- [16] C. K. Burrell and T. J. Osborne. Bounds on the speed of information propagation in disordered quantum spin chains. *Phys. Rev. Lett.*, **99**:167201, (2007).
- [17] M. S. Sarandy. Classical correlation and quantum discord in critical systems. *Phys. Rev. A*, **80**:022108, (2009).
- [18] Z. Yong, L. Gui-Lu, W. Yu-Chun, and G. Guang-Can. Partial teleportation of entanglement through natural thermal entanglement in two-qubit heisenberg xxx chain. *Communications in Theoretical Physics*, **47**:787, (2007).
- [19] Elisa Ercolessi, Stefano Evangelisti, and Francesco Ravanini. Exact entanglement entropy of the xyz model and its sine-gordon limit. *Physics Letters A*, **374**:2101 – 2105, (2010).
- [20] A. Abdel-Aty, L. Y. Cheong, N. Zakaria, and N. Metwally. Entanglement and teleportation via partial entangled-state quantum network. *Journal of Computational and Theoretical Nanoscience*, **12**(9):2213–2220, 2015.
- [21] F. Kheirandish, S. J. Akhtarshenas, and H. Mohammadi. Effect of spin-orbit interaction on entanglement of two-qubit heisenberg xyz systems in an inhomogeneous magnetic field. *Phys. Rev. A*, **77**:042309, (2008).
- [22] X. Xiao-Bo, L. Jin-Ming, and Y. Peng-Fei. Entanglement of a two-qubit anisotropic heisenberg xyz chain in nonuniform magnetic fields with intrinsic decoherence. *Chinese Physics B*, **17**:456, (2008).
- [23] B. S. Shastry. Exact solution of an s=1/2 heisenberg antiferromagnetic chain with long-ranged interactions. *Phys. Rev. Lett.*, **60**:639–642, (1988).
- [24] F. D. M. Haldane. Exact jastrow-gutzwiller resonating-valence-bond ground state of the spin-(1/2) antiferromagnetic heisenberg chain with $1/r^2$ exchange. *Phys. Rev. Lett.*, **60**:635–638, (1988).
- [25] X. S. Ma, Y. Qiao, G. X. Zhao, and A. M. Wang. Quantum discord of thermal states of a spin chain with calogero-moser type interaction. *Science China Physics, Mechanics and Astronomy*, **56**:600–605, (2013).
- [26] M. Avellino, A. J. Fisher, and S. Bose. Quantum communication in spin systems with long-range interactions. *Phys. Rev. A*, **74**:012321, (2006).




Fig. 1: The dynamics of the information (a) Shannon entropy (b) Von Neumann entropy as a function of distance between the Qubits R and the thermalization T.

- [27] S. Bose. Quantum communication through spin chain dynamics: an introductory overview. *Contemporary Physics*, **48**:13–30, (2007).
- [28] G. Gualdi, V. Kostak, I. Marzoli, and P. Tombesi. Perfect state transfer in long-range interacting spin chains. *Phys. Rev. A*, **78**:022325, (2008).
- [29] M. A. Jafarizadeh and R. Sufiani. Perfect state transfer over distance-regular spin networks. *Phys. Rev. A*, **77**:022315, (2008).
- [30] M R Soltani, S MahdaviFar, and M Mahmoudi. Entanglement in a two-spin system with long-range interactions. *Chinese Physics B*, **25**(8):087501, (2016).
- [31] Eloisa Cuestas, Mariano Garagiola, Federico M. Pont, Omar Osenda, and Pablo Serra. Long- and short-range interaction footprints in entanglement entropies of two-particle wigner molecules in 2d quantum traps. *Physics Letters A*, **381**(27):2153 – 2160, (2017).
- [32] Zachary Eldredge, Zhe-Xuan Gong, Jeremy T. Young, Ali Hamed Moosavian, Michael Foss-Feig, and Alexey V. Gorshkov. Fast quantum state transfer and entanglement renormalization using long-range interactions. *Phys. Rev. Lett.*, **119**:170503, (2017).



Ahmad Redwan received his B.Sc. in Mathematics from Al-Azhar University, Assiut, Egypt in 2011 and pre-master courses from Mathematics Department (Applied Mathematics) from Sohag University, Sohag, Egypt, May 2014. He

is currently a demonstrator and master student at Mathematics Department, Faculty of Science, Al-Azhar University, Assiut, Egypt. His research interests include the Information Theory, Quantum Optics and Quantum Information..



Abdel-Haleem Abdel-Aty received his B.Sc. and M.Sc. degrees both in Physics from Al-Azhar University, Egypt, in 2004 and 2008, respectively, and the Ph.D. degree in Computer and Information Science from Universiti Teknologi PETRONAS,

Malaysia, in 2014. He is currently an Assistant Professor at Al-Azhar University and in 2017 he selected as an affiliate with African Academy of Science (AAS). His research interests include the telecommunication, information theory, quantum information, and quantum optics. He published more than 30 papers in peer-reviewed journals and international conferences.



Nour Zidan received his B.Sc and M.Sc degree both in Applied Mathematics from South Valley University, Egypt in 1991 and 1996, respectively, and Ph.D degree in applied Mathematics from Sohag University in 2000. He is currently an Associate Professor at Sohag

University. His research interests include quantum mechanics, quantum information and quantum optics. He published more than 20 papers in peer-reviewed journals and international conferences.



Tarek M. Elshahat

is Associate Professor in the department of Mathematic, Al-Azhar Univrsity. He received the PhD degree in applied Mathematics "Quantum Opitcs" from the Faculty of Science, Al-Azhar University. He was visiting

professor in the group of professor R. Tanas at Adam Mickiewicz University, Poznan, Poland. His research interests are in the areas of applied mathematics, quantum optics, quantum information.