

N -Group SU -Action and its Applications to N -Group Theory

Aslihan Sezgin^{1,*}, Akın Osman Atagün² and Naim Çağman³

¹ Department of Mathematics, Amasya University, 05100 Amasya, Turkey

² Yozgat, Turkey

³ Department of Mathematics, Gaziosmanpaşa University, Tokat, Turkey

Received: 22 Jul. 2016, Revised: 10 Aug. 2016, Accepted: 22 Aug. 2016

Published online: 1 Sep. 2016

Abstract: In this paper, we define a new type of N -group action, called N -group soft union (SU) action on a soft set. This new concept illustrates how a soft set effects on an N -group structure in the mean of union and inclusion of sets and it functions as a bridge among soft set theory, set theory and N -group theory. Furthermore, we derive its basic properties with illustrative examples, investigate the relationship between N -group SI -action defined in [32] and N -group SU -action and obtain some analog of classical N -group theoretic concepts for N -group SU -action. Finally, we give the applications of N -group SU -actions to N -group theory.

Keywords: Soft sets, N -group SI -action, N -group SU -action, N -ideal SU -action, soft pre-image, soft anti image, α -inclusion.

1 Introduction

Molodtsov [23] introduced soft set theory in 1999 for dealing with uncertainties and it continues to experience tremendous growth and diversification in the mean of algebraic structures as in [1, 2, 10, 14, 15, 16, 18, 19, 26, 28, 29, 30, 31, 34].

Operations of soft sets have been studied by some authors. Maji et al. [20] presented some definitions on soft sets and based on the analysis of several operations on soft sets Ali et al. [3] introduced several operations of soft sets and Sezgin and Atagün [27] studied on soft set operations as well. Moreover, soft set relations and functions [4] and soft mappings [22] with many related concepts were discussed. The theory of soft set also has a wide range of applications especially in soft decision making as in the following studies: [5, 6, 13, 21, 24].

Sezgin et al. [32] introduced a new concept to the literature of N -group, called N -group soft intersection action and abbreviated as " N -group SI -action". In this paper, we define a new type of N -group action on a soft set, which we call N -group soft union action and abbreviate as " N -group SU -action". While N -group SI -action is based on the inclusion relation and intersection of sets, N -group SU -action is based on the inclusion relation and union of sets. Since N -group

SU -action gathers soft set theory, set theory and N -group theory, it is useful in improving the soft set theory with respect to N -group structures. Based on this new concept, we then introduce the concepts of N -ideal SU -action and we show that if N is a zero-symmetric near-ring, then every N -ideal SU -action over U is an N -group SU -action over U . Moreover, we investigate these notions with respect to soft pre-image, soft anti image and α -inclusion of soft sets and obtain a significant relationship between N -group SI -action and N -group SU -action. Finally, we give some applications of N -group SU -action to N -group theory.

2 Preliminaries

In this section, we recall some basic notions relevant to N -groups and soft sets. By a *near-ring*, we shall mean an algebraic system $(N, +, \cdot)$, where

N1) $(N, +)$ forms a group (not necessarily abelian)

N2) (N, \cdot) forms a semigroup and

N3) $(a + b)c = ac + bc$ for all $a, b, c \in N$ (i.e. we study on right near-rings.)

Throughout this paper, N will always denote a right near-ring. A normal subgroup I of N is called a left ideal of N

* Corresponding author e-mail: aslihan.sezgin@amasya.edu.tr

if $n(s+i) - ns \in I$ for all $n, s \in N$ and $i \in I$ and denoted by $I \triangleleft_\ell N$.

Let $(\Gamma, +)$ be a group and

$$\begin{aligned} \mu : N \times \Gamma &\rightarrow \Gamma \\ (n, \gamma) &\rightarrow n\gamma \end{aligned}$$

(Γ, μ) is called a *near-ring module* or N -group if $\forall x, y \in N, \forall \gamma \in \Gamma$,

- i) $x(y\gamma) = (xy)\gamma$ and
- ii) $(x+y)\gamma = x\gamma + y\gamma$.

It is denoted by Γ . Clearly N itself is an N -group by natural operation. Let G be a group, written additively but not necessarily abelian, and let $M(G)$ be the set $\{f|f : G \rightarrow G\}$ of all functions from G to G . An addition operation can be defined on $M(G)$: given f, g in $M(G)$, then the mapping $f+g$ from G to G is given by $(f+g)(x) = f(x) + g(x)$ for all x in G . Then $(M(G), +)$ is also a group, which is abelian if and only if G is abelian. Taking the composition of mappings as the product, $M(G)$ becomes a near-ring. Let G be a group. Then, under the operation below:

$$\begin{aligned} \mu : M(G) \times G &\rightarrow G \\ (f, a) &\rightarrow f(a) \end{aligned}$$

G is an $M(G)$ -group. For a near-ring N , the zero-symmetric part of N denoted by N_0 is defined by $N_0 = \{n \in N \mid n0 = 0\}$. A subgroup Δ of Γ with $N\Delta \subseteq \Delta$ is said to be an N -subgroup of Γ and denoted by $\Delta \leq_N \Gamma$. A normal subgroup Δ of Γ is called an N -ideal of Γ and denoted by $\Delta \trianglelefteq_N \Gamma$, if $\forall \gamma \in \Gamma, \forall \delta \in \Delta, \forall n \in N, n(\gamma + \delta) - n\gamma \in \Delta$. Let N be a near-ring, Γ and Ψ two N -groups. Then, $h : \Gamma \rightarrow \Psi$ is called an N -homomorphism if $\forall \gamma, \delta \in \Gamma, \forall n \in N$,

- i) $h(\gamma + \delta) = h(\gamma) + h(\delta)$ and
- ii) $h(n\gamma) = nh(\gamma)$.

For all undefined concepts and notions we refer to [25]. From now on, U refers to an initial universe, E is a set of parameters, $P(U)$ is the power set of U and $A, B, C \subseteq E$.

Definition 1.[6, 23] A soft set f_A over U is a set defined by

$$f_A : E \rightarrow P(U) \text{ such that } f_A(x) = \emptyset \text{ if } x \notin A.$$

Here, f_A is also called *approximate function*. A soft set over U can be represented by the set of ordered pairs

$$f_A = \{(x, f_A(x)) : x \in E, f_A(x) \in P(U)\}.$$

It is clear to see that a soft set is a parametrized family of subsets of the set U . It is worth noting that the sets $f_A(x)$ may be arbitrary. Some of them may be empty, some may have nonempty intersection. If we define more than one soft set in a subset A of the set of parameters E , then the soft sets will be denoted by f_A, g_A, h_A etc. If we define more than one soft set in some subsets A, B, C etc. of parameters E , then the soft sets will be denoted by f_A, f_B, f_C etc., respectively. We refer to [6, 11, 12, 20, 23] for further details.

Definition 2.[6] Let f_A and f_B be soft sets over U . Then, f_A is a soft subset of f_B , denoted by $f_A \subseteq f_B$, if $f_A(x) \subseteq f_B(x)$ for all $x \in E$.

Complement of the soft set f_A over U , denoted by f_A^c , is defined as $f_A^c(\alpha) = U \setminus f_A(\alpha)$ for all $\alpha \in E$.

Definition 3.[6] Let f_A and f_B be soft sets over U . Then, union of f_A and f_B , denoted by $f_A \cup f_B$, is defined as $f_A \cup f_B = f_{A \cup B}$, where $f_{A \cup B}(x) = f_A(x) \cup f_B(x)$ for all $x \in E$.

Intersection of f_A and f_B , denoted by $f_A \cap f_B$, is defined as $f_A \cap f_B = f_{A \cap B}$, where $f_{A \cap B}(x) = f_A(x) \cap f_B(x)$ for all $x \in E$.

Definition 4.[6] Let f_A and f_B be soft sets over U . Then, \vee -product of f_A and f_B , denoted by $f_A \vee f_B$, is defined as $f_A \vee f_B = f_{A \vee B}$, where $f_{A \vee B}(x, y) = f_A(x) \cup f_B(y)$ for all $(x, y) \in E \times E$.

\wedge -product of f_A and f_B , denoted by $f_A \wedge f_B$, is defined as $f_A \wedge f_B = f_{A \wedge B}$, where $f_{A \wedge B}(x, y) = f_A(x) \cap f_B(y)$ for all $(x, y) \in E \times E$.

Definition 5.[7] Let f_A and f_B be soft sets over the common universe U and Ψ be a function from A to B . Then, soft image of f_A under Ψ , denoted by $\Psi(f_A)$, is a soft set over U by

$$(\Psi(f_A))(b) = \begin{cases} \bigcup \{f_A(a) \mid a \in A \text{ and } \Psi(a) = b\}, & \text{if } \Psi^{-1}(b) \neq \emptyset, \\ \emptyset, & \text{otherwise} \end{cases}$$

for all $b \in B$. And soft pre-image (or soft inverse image) of f_B under Ψ , denoted by $\Psi^{-1}(f_B)$, is a soft set over U by $(\Psi^{-1}(f_B))(a) = f_B(\Psi(a))$ for all $a \in A$.

Definition 6.[8] Let f_A and f_B be soft sets over the common universe U and Ψ be a function from A to B . Then, soft anti image of f_A under Ψ , denoted by $\Psi^*(f_A)$, is a soft set over U by $(\Psi^*(f_A))(b) = \begin{cases} \bigcap \{f_A(a) \mid a \in A \text{ and } \Psi(a) = b\}, & \text{if } \Psi^{-1}(b) \neq \emptyset, \\ \emptyset, & \text{otherwise} \end{cases}$ for all $b \in B$.

Theorem 1.[8] Let f_A and f_B be soft sets over U , f_A^c, f_B^c be their complements, respectively and Ψ be a function from A to B . Then,

- i) $\Psi^{-1}(f_B^c) = (\Psi^{-1}(f_B))^c$.
- ii) $\Psi(f_A^c) = (\Psi^*(f_A))^c$ and $\Psi^*(f_A) = (\Psi(f_A))^c$.

Definition 7.[9] Let f_A be a soft set over U and α be a subset of U . Then, upper α -inclusion of f_A , denoted by $f_A^{\supseteq \alpha}$, and lower α -inclusion of f_A , denoted by $f_A^{\subseteq \alpha}$, are defined as

$f_A^{\supseteq \alpha} = \{x \in A \mid f_A(x) \supseteq \alpha\}$ and $f_A^{\subseteq \alpha} = \{x \in A \mid f_A(x) \subseteq \alpha\}$, respectively.

3 N -group SU -actions and N -ideal SU -actions

In this section, we first define N -group soft union actions, abbreviated as N -group SU -actions and N -ideal SU -actions with illustrative examples. We then study their basic properties with respect to soft set operations.

Definition 8. Let Γ be an N -group and f_Γ be a soft set over U . Then, f_Γ is called a N -group SU -action over U if it satisfies the following properties:

- i) $f_\Gamma(x+y) \subseteq f_\Gamma(x) \cup f_\Gamma(y)$,
- ii) $f_\Gamma(-x) = f_\Gamma(x)$,
- iii) $f_\Gamma(nx) \subseteq f_\Gamma(x)$

for all $x, y \in \Gamma$ and $n \in N$.

Example 1. Let $N = \{0, 1, 2, 3\}$ be the (right) near-ring due to [25] (Near-rings of low order (D-5)) with the following tables:

+	0	1	2	3	·	0	1	2	3
0	0	1	2	3	0	0	0	0	0
1	1	2	3	0	1	0	1	1	0
2	2	3	0	1	2	0	2	2	0
3	3	0	1	2	3	0	3	3	0

Let $\Gamma = N$ be the sets of parameters and $U = \left\{ \begin{bmatrix} x & 0 \\ x & 0 \end{bmatrix} \mid x, y \in \mathbb{Z}_4 \right\}$, 2×2 matrices with \mathbb{Z}_4 terms, is the universal set. We construct a soft set f_Γ over U by

$$f_\Gamma(0) = \left\{ \begin{bmatrix} 2 & 0 \\ 2 & 0 \end{bmatrix} \right\}$$

$$f_\Gamma(1) = f_\Gamma(2) = f_\Gamma(3) = \left\{ \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 2 & 0 \end{bmatrix}, \begin{bmatrix} 3 & 0 \\ 3 & 0 \end{bmatrix} \right\}.$$

Then, one can easily show that the soft set f_Γ is an N -group SU -action over U .

Example 2. In Example 1, assume that Γ is again the set of parameters and $U = S_4$ is the universal set. We define a soft set f_Γ by

$$\begin{aligned} f_\Gamma(0) &= \{e\}, & f_\Gamma(1) &= \{e, (13)(24)\}, \\ f_\Gamma(2) &= \{e, (12)(34), (1234), (2134)\} \text{ and} \\ f_\Gamma(3) &= \{e, (13)(24), (134)\}. \end{aligned}$$

Since $f_\Gamma(2 \cdot 1) = f_\Gamma(2) \not\subseteq f_\Gamma(1)$, f_Γ is not an N -group SU -action over U .

It is known that if $N = N_0$, then $n0_\Gamma = 0_\Gamma$ for all $n \in N$. Therefore, if N is a zero-symmetric near-ring and if we take $\Gamma = \{0_\Gamma\}$, then f_Γ is an N -group SU -action over U no matter how f_Γ is defined and no matter what U is.

Proposition 1. Let f_Γ be an N -group SU -action over U . Then, $f_\Gamma(0_\Gamma) \subseteq f_\Gamma(x)$ for all $x \in \Gamma$.

Proof. Assume that f_Γ is an N -group SU -action over U . Then, for all $x \in \Gamma$, $f_\Gamma(0_\Gamma) = f_\Gamma(x - x) \subseteq f_\Gamma(x) \cup f_\Gamma(-x) = f_\Gamma(x) \cup f_\Gamma(x) = f_\Gamma(x)$.

Theorem 2. Let Γ be an N -group and f_Γ be a soft set over U . Then, f_Γ is an N -group SU -action over U if and only if

- i) $f_\Gamma(x-y) \subseteq f_\Gamma(x) \cup f_\Gamma(y)$
- ii) $f_\Gamma(nx) \subseteq f_\Gamma(x)$

for all $x, y \in \Gamma$ and $n \in N$.

Proof. Suppose that f_Γ is an N -group SU -action over. Then, by Definition 8, $f_\Gamma(xy) \subseteq f_\Gamma(y)$ and $f_\Gamma(x-y) \subseteq f_\Gamma(x) \cup f_\Gamma(-y) = f_\Gamma(x) \cup f_\Gamma(y)$ for all $x, y \in \Gamma$.

Conversely, assume that $f_\Gamma(xy) \subseteq f_\Gamma(y)$ and $f_\Gamma(x-y) \subseteq f_\Gamma(x) \cup f_\Gamma(y)$ for all $x, y \in \Gamma$. If we choose $x = 0_\Gamma$, then

$$f_\Gamma(0_\Gamma - y) = f_\Gamma(-y) \subseteq f_\Gamma(0_\Gamma) \cup f_\Gamma(y) = f_\Gamma(y)$$

by Proposition 1. Similarly, $f_\Gamma(y) = f_\Gamma(-(-y)) \subseteq f_\Gamma(-y)$, thus $f_\Gamma(-y) = f_\Gamma(y)$ for all $y \in \Gamma$. Also, by assumption $f_\Gamma(x+y) \subseteq f_\Gamma(x) \cup f_\Gamma(-y) = f_\Gamma(x) \cup f_\Gamma(y)$. Thus, the proof is completed.

Theorem 3. Let f_Γ be an N -group SU -action over U . If $f_\Gamma(x-y) = f_\Gamma(0_\Gamma)$ for any $x, y \in \Gamma$, then $f_\Gamma(x) = f_\Gamma(y)$.

Proof. Assume that $f_\Gamma(x-y) = f_\Gamma(0_\Gamma)$ for any $x, y \in \Gamma$. Then,

$$\begin{aligned} f_\Gamma(x) &= f_\Gamma(x-y+y) \\ &\subseteq f_\Gamma(x-y) \cup f_\Gamma(y) \\ &= f_\Gamma(0_\Gamma) \cup f_\Gamma(y) \\ &= f_\Gamma(y) \end{aligned}$$

and accordingly

$$\begin{aligned} f_\Gamma(y) &= f_\Gamma((y-x)+x) \\ &\subseteq f_\Gamma(y-x) \cup f_\Gamma(x) \\ &= f_\Gamma(-(y-x)) \cup f_\Gamma(x) \\ &= f_\Gamma(0_\Gamma) \cup f_\Gamma(x) \\ &= f_\Gamma(x). \end{aligned}$$

Thus, $f_\Gamma(x) = f_\Gamma(y)$, completing the proof.

It is known that if Γ is an N -group, then $(\Gamma, +)$ is a group but not necessarily abelian. That is, for any $x, y \in \Gamma$, $x+y$ needs not be equal to $y+x$. However, we have the following:

Theorem 4. Let f_Γ be an N -group SU -action over U and $x \in \Gamma$. Then, for all $y \in \Gamma$

$$f_\Gamma(x) = f_\Gamma(0_\Gamma) \Leftrightarrow f_\Gamma(x+y) = f_\Gamma(y+x) = f_\Gamma(y)$$

Proof. Suppose that $f_\Gamma(x+y) = f_\Gamma(y+x) = f_\Gamma(y)$ for all $y \in \Gamma$. Then by choosing $y = 0_\Gamma$, we obtain that $f_\Gamma(x) = f_\Gamma(0_\Gamma)$. Conversely, assume that $f_\Gamma(x) = f_\Gamma(0_\Gamma)$. Then, by Proposition 1, we have

$$f_\Gamma(0_\Gamma) = f_\Gamma(x) \subseteq f_\Gamma(y), \quad \forall y \in \Gamma. \tag{1}$$

Since f_Γ is an N -group SU -action over U , then

$$f_\Gamma(x+y) \subseteq f_\Gamma(x) \cup f_\Gamma(y) = f_\Gamma(y), \quad \forall y \in \Gamma.$$

Furthermore, for all $y \in \Gamma$

$$\begin{aligned} f_\Gamma(y) &= f_\Gamma((-x)+x+y) \\ &= f_\Gamma(-x+(x+y)) \\ &\subseteq f_\Gamma(-x) \cup f_\Gamma(x+y) \\ &= f_\Gamma(x) \cup f_\Gamma(x+y) \\ &= f_\Gamma(x+y) \end{aligned}$$

Because, by (1), $f_{\Gamma}(x) \subseteq f_{\Gamma}(y)$ for all $y \in \Gamma$ and $x, y \in \Gamma$ implies that $x + y \in \Gamma$. Thus, $f_{\Gamma}(x) \subseteq f_{\Gamma}(x + y)$ and it follows that $f_{\Gamma}(x + y) = f_{\Gamma}(y)$ for all $y \in \Gamma$. Now, let $x \in \Gamma$. Then, for all $y \in \Gamma$

$$\begin{aligned} f_{\Gamma}(y + x) &= f_{\Gamma}(y + x + (y - y)) \\ &= f_{\Gamma}(y + (x + y) - y) \\ &\subseteq f_{\Gamma}(y) \cup f_{\Gamma}(x + y) \cup f_{\Gamma}(y) \\ &= f_{\Gamma}(y) \cup f_{\Gamma}(x + y) \\ &= f_{\Gamma}(y), \end{aligned}$$

since $f_{\Gamma}(x + y) = f_{\Gamma}(y)$. Moreover, for all $y \in \Gamma$,

$$\begin{aligned} f_{\Gamma}(y) &= f_{\Gamma}(y + (x - x)) \\ &= f_{\Gamma}((y + x) - x) \\ &\subseteq f_{\Gamma}(y + x) \cup f_{\Gamma}(x) \\ &= f_{\Gamma}(y + x) \end{aligned}$$

by (1). It follows that $f_{\Gamma}(y + x) = f_{\Gamma}(y)$, so $f_{\Gamma}(x + y) = f_{\Gamma}(y + x) = f_{\Gamma}(y)$ for all $y \in \Gamma$.

In [32], Sezgin et al. showed that \wedge -product of two *N*-group *SI*-actions over *U* is an *N*-group *SI*-action. However, we have the following for *N*-group *SU*-actions:

Theorem 5. *If f_{Γ} and f_{Δ} are *N*-group *SU*-actions over *U*, then so is $f_{\Gamma} \vee f_{\Delta}$ over *U*.*

Proof. By Definition 4, let $f_{\Gamma} \vee f_{\Delta} = f_{\Gamma \vee \Delta}$, where $f_{\Gamma \vee \Delta}(x, y) = f_{\Gamma}(x) \cup f_{\Delta}(y)$ for all $(x, y) \in E \times E$. Since Γ and Δ are *N*-groups, then $\Gamma \times \Delta$ is an $N \times N$ -group. So, let $(x_1, y_1), (x_2, y_2) \in \Gamma \times \Delta$ and $(n_1, n_2) \in N \times N$. Then,

$$\begin{aligned} f_{\Gamma \vee \Delta}((x_1, y_1) - (x_2, y_2)) &= f_{\Gamma \vee \Delta}(x_1 - x_2, y_1 - y_2) \\ &= f_{\Gamma}(x_1 - x_2) \cup f_{\Delta}(y_1 - y_2) \\ &\subseteq (f_{\Gamma}(x_1) \cup f_{\Gamma}(x_2)) \cup (f_{\Delta}(y_1) \cup f_{\Delta}(y_2)) \\ &= (f_{\Gamma}(x_1) \cup f_{\Delta}(y_1)) \cup (f_{\Gamma}(x_2) \cup f_{\Delta}(y_2)) \\ &= f_{\Gamma \vee \Delta}(x_1, y_1) \cup f_{\Gamma \vee \Delta}(x_2, y_2) \end{aligned}$$

$$\begin{aligned} f_{\Gamma \vee \Delta}((n_1, n_2)(x_1, y_1)) &= f_{\Gamma \vee \Delta}(n_1 x_1, n_2 y_1) \\ &= f_{\Gamma}(n_1 x_1) \cup f_{\Delta}(n_2 y_1) \\ &\subseteq f_{\Gamma}(x_1) \cup f_{\Delta}(y_1) \\ &= f_{\Gamma \vee \Delta}(x_1, y_1) \end{aligned}$$

Thus, $f_{\Gamma} \vee f_{\Delta}$ is an *N*-group *SU*-action over *U*.

In [32], Sezgin et al. showed that if f_{Γ} and h_{Γ} are two *N*-group *SI*-actions over *U*, then so is $f_{\Gamma} \widetilde{\cup} h_{\Gamma}$ over *U*. However, we have the following for *N*-group *SU*-actions:

Theorem 6. *If f_{Γ} and h_{Γ} are two *N*-group *SU*-actions over *U*, then so is $f_{\Gamma} \widetilde{\cup} h_{\Gamma}$ over *U*.*

Proof. Let $x, y \in \Gamma$ and $n \in N$, then

$$\begin{aligned} (f_{\Gamma} \widetilde{\cup} h_{\Gamma})(x - y) &= f_{\Gamma}(x - y) \cup h_{\Gamma}(x - y) \\ &\subseteq (f_{\Gamma}(x) \cup f_{\Gamma}(y)) \cup (h_{\Gamma}(x) \cup h_{\Gamma}(y)) \\ &= (f_{\Gamma}(x) \cup h_{\Gamma}(x)) \cup (f_{\Gamma}(y) \cup h_{\Gamma}(y)) \\ &= (f_{\Gamma} \widetilde{\cup} h_{\Gamma})(x) \cup (f_{\Gamma} \widetilde{\cup} h_{\Gamma})(y), \end{aligned}$$

$$\begin{aligned} (f_{\Gamma} \widetilde{\cup} h_{\Gamma})(nx) &= f_{\Gamma}(nx) \cup h_{\Gamma}(nx) \\ &\subseteq f_{\Gamma}(x) \cup h_{\Gamma}(x) \\ &= (f_{\Gamma} \widetilde{\cup} h_{\Gamma})(x) \end{aligned}$$

Therefore, $f_{\Gamma} \widetilde{\cup} h_{\Gamma}$ is an *N*-group *SU*-action over *U*.

Definition 9. *Let Γ be an *N*-group and f_{Γ} be an *N*-group *SU*-action over *U*. Then, f_{Γ} is called an *N*-ideal *SU*-action of Γ over *U* if it satisfies the following properties:*

- i) $f_{\Gamma}(x + y) \subseteq f_{\Gamma}(x) \cup f_{\Gamma}(y)$,
- ii) $f_{\Gamma}(-x) = f_{\Gamma}(x)$,
- iii) $f_{\Gamma}(x + y - x) \subseteq f_{\Gamma}(y)$,
- iv) $f_{\Gamma}(n(x + y) - nx) \subseteq f_{\Gamma}(y)$,

for all $x, y \in \Gamma$ and $n \in N$. Here, note that $f_{\Gamma}(x + y) \subseteq f_{\Gamma}(x) \cup f_{\Gamma}(y)$ and $f_{\Gamma}(-x) = f_{\Gamma}(x)$ imply $f_{\Gamma}(x - y) \subseteq f_{\Gamma}(x) \cup f_{\Gamma}(y)$.

Example 3. Let $N = \{0, 1, 2, 3\}$ be the (right) near-ring due to [25] (Near-rings of low order (*D*-10)) with the following tables:

+	0	1	2	3	.	0	1	2	3
0	0	1	2	3	0	0	0	0	0
1	1	2	3	0	1	0	1	2	1
2	2	3	0	1	2	0	2	0	2
3	3	0	1	2	3	0	3	2	3

Let $\Gamma = N$ be the sets of parameters and $U = D_3$, dihedral group, be the universal set. We define a soft set f_{Γ} over *U* by

$$f_{\Gamma}(0) = \{e, x\}, f_{\Gamma}(1) = f_{\Gamma}(3) = \{e, x, yx, yx^2\}, f_{\Gamma}(2) = \{e, x, yx^2\}.$$

Then, one can show that f_{Γ} is an *N*-ideal *SU*-action of Γ over *U*.

Example 4. Let $N = \{0, a, b, c\}$ be the (right) near-ring per scheme 2 ([25], p. 408) under the operations defined by the following tables:

+	0	a	b	c	.	0	a	b	c
0	0	a	b	c	0	0	0	0	0
a	a	0	c	b	a	0	0	a	a
b	b	c	0	a	b	0	a	b	b
c	c	b	a	0	c	0	a	c	c

Let $\Gamma = N$ be the sets of parameters and $U = \mathbb{Z}^-$ be the universal set. We define a soft set f_{Γ} over *U* by $f_{\Gamma}(0) = \{-3\}$, $f_{\Gamma}(a) = \{-3, -5, -9\}$, $f_{\Gamma}(b) = \{-3, -5, -9, -11, -15\}$,

$$f_{\Gamma}(c) = \{-3, -11, -15\}.$$

Since $f_{\Gamma}(a(c + c) - ac) = f_{\Gamma}(a0 - ac) = f_{\Gamma}(0 - a) = f_{\Gamma}(0 + a) = f_{\Gamma}(a) \not\subseteq f_{\Gamma}(c)$, f_{Γ} is not an *N*-ideal *SU*-action of Γ over *U*.

It is known that if *N* is a zero-symmetric near-ring, then every *N*-ideal of Γ is also an *N*-subgroup of Γ [25]. Here, we have an analog for this case:

Theorem 7. *Let *N* be a zero-symmetric near-ring. Then, every *N*-ideal *SU*-action over *U* is an *N*-group *SU*-action over *U*.*

Proof. Let f_Γ be an N -ideal SU -action of Γ over U . Since $f_\Gamma(n(x+y) - nx) \subseteq f_\Gamma(y)$, for all $x, y \in \Gamma$ and $n \in N$, in particular for $x = 0_\Gamma$, it follows that $f_\Gamma(n(0_\Gamma + y) - n0_\Gamma) = f_\Gamma(ny - 0_\Gamma) = f_\Gamma(ny) \subseteq f_\Gamma(y)$. Since the other conditions is satisfied by Definition 9, f_Γ is an N -group SU -action over U .

In [32], Sezgin et al. showed that \wedge -product of two N -ideal SI -actions over U is an N -ideal SI -action over U . However, we have the following for N -ideal SU -action:

Theorem 8. *If f_Γ is an N -ideal SU -action of Γ and f_Δ is an N -ideal SU -action of Δ over U , then $f_\Gamma \vee f_\Delta$ is an N -ideal SU -action of $\Gamma \times \Delta$ over U .*

Proof. Let $(x_1, y_1), (x_2, y_2)$ and $(n_1, n_2) \in N \times N$. Then $f_{\Gamma \vee \Delta}((x_1, y_1) - (x_2, y_2)) \subseteq f_{\Gamma \vee \Delta}(x_1, y_1) \cup f_{\Gamma \vee \Delta}(x_2, y_2)$ can be shown similar to Theorem 5. Now,

$$\begin{aligned} f_{\Gamma \vee \Delta}((x_1, y_1) + (x_2, y_2) - (x_1, y_1)) &= f_{\Gamma \vee \Delta}(x_1 + x_2 - x_1, y_1 + y_2 - y_1) \\ &= f_\Gamma(x_1 + x_2 - x_1) \cup f_\Delta(y_1 + y_2 - y_1) \\ &\subseteq f_\Gamma(x_2) \cup f_\Delta(y_2) \\ &= f_{\Gamma \vee \Delta}(x_2, y_2), \end{aligned}$$

and

$$\begin{aligned} f_{\Gamma \vee \Delta}((n_1, n_2)((x_1, y_1) + (x_2, y_2)) - (n_1, n_2)(x_1, y_1)) \\ &= f_{\Gamma \vee \Delta}(n_1(x_1 + x_2) - n_1x_1, n_2(y_1 + y_2) - n_2y_1) \\ &= f_\Gamma(n_1(x_1 + x_2) - n_1x_1) \cup f_\Delta(n_2(y_1 + y_2) - n_2y_1) \\ &\subseteq f_\Gamma(x_2) \cup f_\Delta(y_2) \\ &= f_{\Gamma \vee \Delta}(x_2, y_2). \end{aligned}$$

Therefore, $f_\Gamma \vee f_\Delta$ is an N -ideal SU -action of $\Gamma \times \Delta$ over U .

In [32], Sezgin et al. showed that if f_Γ and h_Γ are two N -ideal SI -actions of Γ over U , then so is $f_\Gamma \tilde{\cap} h_\Gamma$ over U . However, we have the following for N -ideal SU -actions:

Theorem 9. *If f_Γ and h_Γ are two N -ideal SU -actions of Γ over U , then $f_\Gamma \tilde{\cup} h_\Gamma$ is an N -ideal SU -action of Γ over U .*

Proof. Let $x, y \in \Gamma$ and $n \in N$. Then,

$$(f_\Gamma \tilde{\cup} h_\Gamma)(x - y) \subseteq (f_\Gamma \tilde{\cup} h_\Gamma)(x) \cup (f_\Gamma \tilde{\cup} h_\Gamma)(y)$$

can be shown similar to Theorem 6. Now,

$$\begin{aligned} (f_\Gamma \tilde{\cup} h_\Gamma)(x + y - x) &= f_\Gamma(x + y - x) \cup h_\Gamma(x + y - x) \\ &\subseteq f_\Gamma(y) \cup h_\Gamma(y) \\ &= (f_\Gamma \tilde{\cup} h_\Gamma)(y) \end{aligned}$$

$$\begin{aligned} (f_\Gamma \tilde{\cup} h_\Gamma)(n(x + y) - nx) &= f_\Gamma(n(x + y) - nx) \cup h_\Gamma(n(x + y) - nx) \\ &\subseteq f_\Gamma(ny) \cup h_\Gamma(ny) \\ &= (f_\Gamma \tilde{\cup} h_\Gamma)(y) \end{aligned}$$

Therefore, $f_\Gamma \tilde{\cup} h_\Gamma$ is an N -ideal SU -action of Γ over U .

4 Applications of N -group SU -actions and N -ideal SU -actions

In this section, first we obtain the relation between N -ideal SI -action and N -ideal SU -action of an N -group over U and then give the applications of soft pre-image, soft anti image, lower α -inclusion of soft sets and N -homomorphism to N -group theory with respect to N -group SU -actions and N -ideal SU -actions.

Theorem 10. *Let f_Γ be a soft set over U . Then, f_Γ is an N -ideal SU -action of Γ over U if and only if f_Γ^c is an N -ideal SI -action of Γ over U .*

Proof. Let f_Γ be an N -ideal SU -action of Γ over U . Then, for all $x, y \in \Gamma$ and $n \in N$,

$$\begin{aligned} f_\Gamma^c(x - y) &= U \setminus f_\Gamma(x - y) \\ &\supseteq U \setminus ((f_\Gamma(x) \cup f_\Gamma(y))) \\ &= (U \setminus f_\Gamma(x)) \cap (U \setminus f_\Gamma(y)) \\ &= f_\Gamma^c(x) \cap f_\Gamma^c(y), \end{aligned}$$

Also,

$$\begin{aligned} f_\Gamma^c(x + y - x) &= U \setminus f_\Gamma(x + y - x) \\ &\supseteq U \setminus (f_\Gamma(y)) \\ &= f_\Gamma^c(y) \end{aligned}$$

Furthermore,

$$\begin{aligned} f_\Gamma^c(n(x + y) - nx) &= U \setminus f_\Gamma(n(x + y) - nx) \\ &\supseteq U \setminus (f_\Gamma(y)) \\ &= f_\Gamma^c(y) \end{aligned}$$

which shows that f_Γ^c is an N -ideal SI -action of Γ over U . The converse can be shown similarly.

Theorem 11. *If f_Γ is an N -ideal SU -action of Γ over U , then $\Gamma_f = \{x \in \Gamma : f_\Gamma(x) = f_\Gamma(0_\Gamma)\}$ is an N -ideal of Γ .*

Proof. It is obvious that $0_\Gamma \in \Gamma_f \subseteq \Gamma$. We need to show that (i) $x - y \in \Gamma_f$, (ii) $\gamma + x - \gamma \in \Gamma_f$ and (iii) $n(\gamma + x) - n\gamma \in \Gamma_f$ for all $x, y \in \Gamma_f$ and $n \in N$ and $\gamma \in \Gamma$. If $x, y \in \Gamma_f$, then $f_\Gamma(x) = f_\Gamma(y) = f_\Gamma(0_\Gamma)$. By Proposition 1,

$f_\Gamma(0_\Gamma) \subseteq f_\Gamma(x - y)$, $f_\Gamma(0_\Gamma) \subseteq f_\Gamma(\gamma + x - \gamma)$ and $f_\Gamma(0_\Gamma) \subseteq f_\Gamma(n(\gamma + x) - n\gamma)$ for all $n \in N$, $x, y \in \Gamma_f$ and $\gamma \in \Gamma$. Since f_Γ is an N -ideal SU -action of Γ over U , then for all $n \in N$, $x, y \in \Gamma_f$ and $\gamma \in \Gamma$

- (i) $f_\Gamma(x - y) \subseteq f_\Gamma(x) \cup f_\Gamma(y) = f_\Gamma(0_\Gamma)$,
- (ii) $f_\Gamma(\gamma + x - \gamma) \subseteq f_\Gamma(x) = f_\Gamma(0_\Gamma)$ and
- (iii) $f_\Gamma(n(\gamma + x) - n\gamma) \subseteq f_\Gamma(x) = f_\Gamma(0_\Gamma)$.

Hence,

$f_\Gamma(x - y) = f_\Gamma(0_\Gamma)$, $f_\Gamma(\gamma + x - \gamma) = f_\Gamma(0_\Gamma)$ and $f_\Gamma(n(\gamma + x) - n\gamma) = f_\Gamma(0_\Gamma)$ for all $n \in N$, $x, y \in \Gamma_f$ and $\gamma \in \Gamma$. Therefore, Γ_f is an N -ideal of Γ .

Theorem 12. [32] *Let f_Γ be a soft set over U and α be a subset of U such that $\emptyset \subseteq \alpha \subseteq f_\Gamma(0_\Gamma)$. If f_Γ is an N -ideal SI -action over U , then $f_\Gamma^{\subseteq \alpha}$ is an N -ideal of Γ .*

Theorem 13. *Let f_Γ be a soft set over U and α be a subset of U such that $\emptyset \subseteq f_\Gamma(0_\Gamma) \subseteq \alpha$. If f_Γ is an N -ideal SU -action of Γ over U , then $f_\Gamma^{\subseteq \alpha}$ is an ideal of Γ .*

Proof. Since $f_\Gamma(0_\Gamma) \subseteq \alpha$, then $0_\Gamma \in f_\Gamma^{\subseteq \alpha}$ and $\emptyset \neq f_\Gamma^{\subseteq \alpha} \subseteq \Gamma$. Let $x, y \in f_\Gamma^{\subseteq \alpha}$, then

$$f_\Gamma(x) \subseteq \alpha \text{ and } f_\Gamma(y) \subseteq \alpha.$$

We need to show that (i) $x - y \in f_{\Gamma}^{\subseteq \alpha}$, (ii) $\gamma + x - \gamma \in f_{\Gamma}^{\subseteq \alpha}$ and (iii) $n(\gamma + x) - n\gamma \in f_{\Gamma}^{\subseteq \alpha}$ for all $x, y \in f_{\Gamma}^{\subseteq \alpha}$, $n \in N$ and $\gamma \in \Gamma$. Since f_{Γ} is an N -ideal SI -action of Γ over U , it follows that

$$\begin{aligned} f_{\Gamma}(x - y) &\subseteq f_{\Gamma}(x) \cup f_{\Gamma}(y) \subseteq \alpha \cup \alpha = \alpha, \\ f_{\Gamma}(\gamma + x - \gamma) &\subseteq f_{\Gamma}(x) \subseteq \alpha \text{ and} \\ f_{\Gamma}(n(\gamma + x) - n) &\subseteq f_{\Gamma}(x) \subseteq \alpha. \end{aligned}$$

Thus, the proof is completed.

Theorem 14.[32] Let f_{Γ} and f_{Δ} be soft sets over U and Ψ be an N -isomorphism from Γ to Δ . If f_{Γ} is an N -ideal SI -action of Γ over U , then $\Psi(f_{\Gamma})$ is an N -ideal SI -action of Δ over U .

Theorem 15. Let f_{Γ} and f_{Δ} be soft sets over U and Ψ be an N -isomorphism from Γ to Δ . If f_{Γ} is an N -ideal SU -action of Γ over U , then $\Psi^*(f_{\Gamma})$ is an N -ideal SU -action of Δ over U .

Proof. Let f_{Γ} be an N -ideal SU -action of Γ over U . Then, f_{Γ}^c is an N -ideal SI -action of Γ over U by Theorem 10 and $\Psi(f_{\Gamma}^c)$ is an N -ideal SI -action of Δ over U by Theorem 14. Thus, $\Psi(f_{\Gamma}^c) = (\Psi^*(f_{\Gamma}))^c$ is an N -ideal SI -action of Δ over U by Theorem 1 (ii). Therefore, $\Psi^*(f_{\Gamma})$ is an N -ideal SU -action of Δ over U by Theorem 10.

Theorem 16.[32] Let f_{Γ} and f_{Δ} be soft sets over U and Ψ be an N -homomorphism from N to Δ . If f_{Δ} is an N -ideal SI -action of Δ over U , then $\Psi^{-1}(f_{\Delta})$ is an N -ideal SI -action of Γ over U .

Theorem 17. Let f_{Γ} and f_{Δ} be soft sets over U and Ψ be an N -homomorphism from Γ to Δ . If f_{Δ} is an N -ideal SU -action of Δ over U , then $\Psi^{-1}(f_{\Delta})$ is an N -ideal SU -action of Γ over U .

Proof. Let f_{Δ} be an N -ideal SU -action of Δ over U . Then, f_{Δ}^c is an N -ideal SI -action of Δ over U by Theorem 10 and $\Psi^{-1}(f_{\Delta}^c)$ is an N -ideal SI -action of Γ over U by Theorem 16. Thus, $\Psi^{-1}(f_{\Delta}^c) = (\Psi^{-1}(f_{\Delta}))^c$ is an N -ideal SI -action of Γ over U by Theorem 1 (i). Therefore, $\Psi^{-1}(f_{\Delta})$ is an N -ideal SU -action of Γ over U by Theorem 10.

5 Conclusion

In this paper, we have defined a new kind of N -group action on a soft set, called N -group SU -action. This new concept is very functional for obtaining results in the mean of N -group structure, since it brings the soft sets, sets and N -groups together. Based on the definition, we have introduced the concept of N -ideal SU -action of an N -group. We have then investigated this notion with respect to soft pre-image, soft anti image and lower α -inclusion of soft sets. Finally, we obtain the relationship between N -group SI -action and N -group SU -action and give some applications of these new concepts to N -group theory. To extend this study, one can further study the other algebraic structures such as algebras in view of their SU -actions.

References

- [1] U. Acar, F. Koyuncu and B. Tanay, *Soft sets and soft rings*, Comput. Math. Appl. **59** (2010), 3458–3463.
- [2] H. Aktaş and N. Çağman, *Soft sets and soft groups*, Inform. Sci. **177** (2007), 2726–2735.
- [3] M.I. Ali, F. Feng, X. Liu, W.K. Min and M. Shabir, *On some new operations in soft set theory*, Comput. Math. Appl. **57** (2009), 1547–1553.
- [4] K.V. Babitha and J.J. Sunil, *Soft set relations and functions*, Comput. Math. Appl. **60** (7) (2010), 1840–1849.
- [5] N. Çağman and S. Enginoğlu, *Soft matrix theory and its decision making*, Comput. Math. Appl. **59** (2010), 3308–3314.
- [6] N. Çağman and S. Enginoğlu, *Soft set theory and uni-int decision making*, Eur. J. Oper. Res. **207** (2010), 848–855.
- [7] N. Çağman, F. Çıtak and H. Aktaş, *Soft int-groups and its applications to group theory*, Neural Comput. Appl., **21** (Issue 1-Supplement) (2012), 151–158.
- [8] N. Çağman, A. Sezgin and A.O. Atagün, *Group SU -action and its applications*, submitted.
- [9] N. Çağman, A. Sezgin and A.O. Atagün, *α -inclusions and their applications to group theory*, submitted.
- [10] F. Feng, Y. B. Jun and X. Zhao, *Soft semirings*, Comput. Math. Appl. **56** (2008), 2621–2628.
- [11] F. Feng, X.Y. Liu, V. Leoreanu-Fotea, Y.B. Jun, *Soft sets and soft rough sets*, Inform. Sci. **181** (6) (2011), 1125–1137.
- [12] F. Feng, C. Li, B. Davvaz and M. I. Ali, *Soft sets combined with fuzzy sets and rough sets: a tentative approach*, Soft Comput. **14** (6) (2010), 899–911.
- [13] F. Feng, Y.M. Li, N. Çağman, *Generalized uni-int decision making schemes based on choice value soft sets*, Eur. J. Oper. Res. **220** (2012), 162–170.
- [14] Y.B. Jun, *Soft BCK/BCI-algebras*, Comput. Math. Appl. **56** (2008), 1408–1413.
- [15] Y.B. Jun and C.H. Park, *Applications of soft sets in ideal theory of BCK/BCI-algebras*, Inform. Sci. **178** (2008), 2466–2475.
- [16] Y.B. Jun, K.J. Lee and J. Zhan, *Soft p -ideals of soft BCI-algebras*, Comput. Math. Appl. **58** (2009), 2060–2068.
- [17] Y.B. Jun, K.J. Lee, C.H. Park, *Soft set theory applied to ideals in d -algebras*, Comput. Math. Appl. **57** (3) (2009), 367–378.
- [18] Y. B. Jun, S. Z. Song and K. S. So, *Soft set theory applied to p -ideals of BCI-algebras related to fuzzy points*, Neural. Comput. Applic. **20** (2011), 1313–1320.
- [19] O. Kazanci, Ş. Yılmaz and S. Yamak, *Soft sets and soft BCH-algebras*, Hacet. J. Math. Stat. **39** (2) (2010), 205–217.
- [20] P.K. Maji, R. Biswas and A.R. Roy, *Soft set theory*, Comput. Math. Appl. **45** (2003), 555–562.
- [21] P.K. Maji, A.R. Roy and R. Biswas, *An application of soft sets in a decision making problem*, Comput. Math. Appl. **44** (2002), 1077–1083.
- [22] P. Majumdar and S.K. Samanta, *On soft mappings*, Comput. Math. Appl. **60** (9) (2010), 2666–2672.
- [23] D. Molodtsov, *Soft set theory-first results*, Comput. Math. Appl. **37** (1999), 19–31.
- [24] D.A. Molodtsov, V. Yu. Leonov and D. V. Kovkov, *Soft sets technique and its application*, Nechetkie Sistemy i Myagkie Vychisleniya **1** (1) (2006), 8–39.
- [25] G. Pilz, *Near-rings*, North Holland Publishing Company, Amsterdam-New York-Oxford, 1983.

- [26] A. Sezgin, A.O. Atagün and E. Aygün, *A note on soft near-rings and idealistic soft near-rings*, *Filomat*, **25** (2011), 53-68.
- [27] A. Sezgin and A.O. Atagün, *On operations of soft sets*, *Comput. Math. Appl.* **61** (5) (2011), 1457–1467.
- [28] A. Sezgin, A.O. Atagün, *Soft groups and normalistic soft groups*, *Comput. Math. Appl.* **62** (2) (2011), 685-698.
- [29] A. Sezgin, A.O. Atagün and N. Çağman, *Union soft substructures of near-rings and N-groups*, *Neural Comput. Appl.* **21** (Issue 1-Supplement) (2012), 133-143.
- [30] A. Sezgin, A.O. Atagün and N. Çağman, *Soft intersection near-rings with applications*, *Neural Comput. Appl.*, **21** (Issue 1-Supplement) (2012), 221-229.
- [31] A. Sezgin Sezer, *A new view to ring theory via soft union rings, ideals and bi-ideals*, *Knowledge-Based Systems*, **36** (2012), 300-314.
- [32] A. Sezgin Sezer, A.O. Atagün and N. Çağman, *N-group SI-action and its applications to N-Group Theory*, *Fasciculi Mathematici*, **54** (2014).
- [33] G. Wendt, *On Zero Divisors in Near-Rings*, *Int. J. Algebra*, **3** (1) (2009), 21–32.
- [34] J. Zhan, Y.B. Jun, *Soft BL-algebras based on fuzzy sets*, *Comput. Math. Appl.* **59** (6) (2010), 2037–2046.



near-ring theory.

Aslihan Sezgin is an Associate Professor Doctor of Mathematics at Amasya University. She received the PhD degree in Mathematics at Gaziosmanpaşa University (Turkey). Her research interests are in the areas of soft set and its algebraic applications and general



Akın Osman Atagün is an Associate Professor of Mathematics. He received the PhD degree in Mathematics at Erciyes University (Turkey). His main research interests are: Near-ring, Prime ideals of near-rings and near-ring modules, Soft set and its algebraic applications.



Education.

Naim Çağman is a Professor of Mathematics at Gaziosmanpaşa University. He received the PhD degree in Mathematics at University of Leeds (UK). His main research interests are: Mathematical Logic, Fuzzy Logic, Soft Set Theory, Game Development, Mathematics