

Four-Parameter Potential Function with Negative Energy Bound States

T. J. Taiwo*

Department of Mathematics, University of Benin, Benin City, Edo State 300283, Nigeria.

Received: 10 Oct. 2018, Revised: 28 Oct. 2018, Accepted: 15 Nov. 2018

Published online: 1 Jan. 2019

Abstract: Using the Tridiagonal Representation Approach (A method where we work in a complete set of square integrable basis that carries a tridiagonal matrix representation for the wave operator. Consequently, the matrix wave equation becomes a three-term recursion relation for the expansion coefficients of the wavefunction. Finding a solution of this recursion relation in terms of orthogonal polynomials is equivalent to solving the original problem) we obtain solutions for a new four-parameter one-dimensional potential function. We obtained the energy spectrum and corresponding wavefunction.

PACS numbers: 03.65.Ge, 03.65.Fd, 34.80.Bm, 03.65.Ca

Keywords: basis set, energy spectrum, wavefunction, tridiagonal representations, negative energy bound states, recursion relation.

1 Introduction and Formulation

Recently, we solved a new five - parameter potential box in one dimensional [1] using TRA. We got the energy spectrum and the wavefunction of this five parameter potential function. In this article we further consider a new four parameter potential function in one dimensional. This potential function is given as:

$$V(x) = \frac{1}{e^{\lambda x} - 1} \left[V_0 + V_1 (1 - 2e^{-\lambda x}) + \frac{V_R}{1 - e^{-\lambda x}} \right] \quad (1)$$

for $0 \leq x \leq \infty$, V_i are real parameters such that V_R is greater than or equal to zero. Fig 1 is the plot of the potential function where V_R is vary (keeping V_0 and V_1 constant) and varying V_0 and V_1 (keeping V_R constant). This potential has never been studied in the published literature. However, when deformed, this potential can be compared to a three - parameter potential in

$$V(x) = V_1 \frac{e^{-\lambda x} - \gamma}{e^{\lambda x} - 1} \quad (2)$$

Where V_1 are the potential strength and the range parameter λ is positive with inverse length unit. The dimensionless parameter γ is in the open range $0 < \gamma < 1$. Graphically, this potential function cross the x -axis at $x_0 = -\ln \gamma / \lambda$ and then reaches a local extremum value of $V(x_1) = -V_0 (1 - \sqrt{1 - \gamma})^2$ at $x_1 = -\frac{1}{\lambda} \ln(1 - \sqrt{1 - \gamma})$. We observed that (1) and (2) are short range potentials with $1/x$ singularity at the origin. Simple algebraic manipulation in (1) will easily produce (2). Note (1) is same as

$$\begin{aligned} V(x) &= \frac{V_0}{e^{\lambda x} - 1} + \frac{V_1(1 - 2e^{-\lambda x})}{e^{\lambda x} - 1} + \frac{V_R e^{\lambda x}}{(e^{\lambda x} - 1)^2} \\ &= V_1 \frac{\gamma - e^{-\lambda x}}{e^{\lambda x} - 1} + \frac{V_R e^{\lambda x}}{(e^{\lambda x} - 1)^2} \end{aligned} \quad (3)$$

When $V_R = 0$, such that we defined a dimensionless ratio

$$\gamma = \frac{V_0 + V_1}{2V_1} \text{ for } V_1 \neq 0; \text{ then we easily get (2).}$$

*Corresponding author e-mail: tundetaiwo31@gmail.com

Detailed study of these potentials; show that they can be used practically as an appropriate model for the interaction of an electron with extended molecules whose electron cloud is congregated near the centre of the molecules. Also, there is a resemblance between this potential and attractive Coulomb potential for nonzero angular momentum at short distances hence we expect them to finite bound states. In this paper, we are more interested in the generalized potential given in (1) where none of the potential parameters is zero in order to obtain the energy spectrum and wave function.

We now formulate this problem and solve it using the Tridiagonal Representation Approach (TRA) [2-7].

We write the wavefunction as the bounded sum $\psi(E, x) = \sum_n f_n(E) \phi_n(x)$, where $\{\phi_n(x)\}$ is a complete set of square integrable basis functions and $\{f_n(E)\}$ are proper expansion coefficients in the energy. If we make the coordinate transformation $y(x) = 1 - 2e^{-\lambda x}$, then the Schrödinger wave equation in the new configuration space becomes

$$(H - E)|\psi\rangle = -\frac{1}{2} \left[(y')^2 \frac{d^2}{dy^2} + y'' \frac{d}{dy} - 2V(y) + 2E \right] |\psi\rangle = 0 \quad (4)$$

where the prime stands for the derivative with respect to x and we adopted the atomic units $\hbar = m = 1$. With $y \in [-1, +1]$, we can choose the following square integrable functions in the new configuration space with coordinate y as basic elements for the expansion of the wavefunction

$$\phi_n(y) = A_n (1-y)^\alpha (1+y)^\beta P_n^{(\mu, \nu)}(y) \quad (5)$$

Where $P_n^{(\mu, \nu)}(x)$ is the Jacobi polynomial of degree $n = 0, 1, 2, \dots$ in y , the parameters μ and ν are larger than -1 and $A_n = \sqrt{\frac{2n+\mu+\nu+1}{2^{\mu+\nu+1}} \frac{\Gamma(n+1)\Gamma(n+\mu+\nu+1)}{\Gamma(n+\nu+1)\Gamma(n+\mu+1)}}$. For simplicity, we rewrite (4) as

$$(H - E)|\psi\rangle = -\frac{1}{2} \left(\frac{(y')^2}{(1-y)^2} \frac{d^2}{dy^2} + \frac{y''}{(y')^2} (1-y^2) \frac{d}{dy} + \frac{2(1-y^2)}{(y')^2} [E - V] \right) |\psi\rangle = 0 \quad (6)$$

We applied the wave operator on the basis element to get

$$(H - E)|\phi_n\rangle = J|\phi_n\rangle = -\frac{\lambda^2 (1-y)}{2(1+y)} \left(\frac{(1-y^2)}{dy^2} - (1+y) \frac{d}{dy} + \frac{2(1+y)E}{\lambda^2(1-y)} - \frac{2V_0}{\lambda^2} - \frac{2V_1 y}{\lambda^2} - \frac{4V_R}{\lambda^2(1+y)} \right) |\phi_n\rangle = 0 \quad (7)$$

where we had used $y' = \lambda(1-y)$, $\frac{y''}{(y')^2} = -1/(1-y)$, and

the potential function in new configuration space as $V(y) = \frac{(1-y)}{(1+y)} \left[V_0 + V_1 y + \frac{2V_R}{(1+y)} \right]$. Using $u_i = \left(\frac{2}{\lambda^2} \right) V_i$ and $\varepsilon = \left(\frac{2}{\lambda^2} \right) E$ in (7) we have

$$\frac{2}{\lambda^2} J|\phi_n\rangle = -\frac{(1-y)}{(1+y)} \left(\frac{(1-y^2)}{dy^2} - (1+y) \frac{d}{dy} - (\varepsilon + u_0) - u_1 y + \frac{2\varepsilon}{(1-y)} - \frac{2u_R}{(1+y)} \right) |\phi_n\rangle = 0 \quad (8)$$

Now, the boundary conditions and square integrability (with respect to the integral measure dx) dictate that the matrix wave operator becomes

$$\begin{aligned} \langle \phi_n | F[y] | \phi_m \rangle &= -A_n A_m \int_{-1}^{+1} (1-y)^{2\alpha} (1+y)^{2\beta} \frac{(1-y)}{(1+y)} [F(y)] P_n^{(\mu, \nu)}(y) P_m^{(\mu, \nu)}(y) \frac{dy}{\lambda(1-y)} \\ &= -\frac{A_n A_m}{\lambda} \int_{-1}^{+1} (1-y)^{2\alpha} (1+y)^{2\beta-1} [F(y)] P_n^{(\mu, \nu)}(y) P_m^{(\mu, \nu)}(y) dy \quad (9) \end{aligned}$$

where

$$F(y) = (1-y^2) \frac{d^2}{dy^2} - (1+y) \frac{d}{dy} - (\varepsilon + u_0) - u_1 y + \frac{2\varepsilon}{(1-y)} - \frac{2u_R}{(1+y)}$$

Hence we have $2\alpha = \mu$ and $2\beta = \nu + 1$. As a result of these conditions with the fact that $y' = \lambda(1-y)$, the normalization constant will be $A_n = \sqrt{\frac{2n+\mu+\nu+1}{2^{\mu+\nu+1}} \frac{\Gamma(n+1)\Gamma(n+\mu+\nu+1)}{\Gamma(n+\nu+1)\Gamma(n+\mu+1)}}$. Using the first and second derivatives of the basis element with respect to y in (8) gives

$$\frac{2}{\lambda^2} J|\phi_n\rangle = -A_n (1-y)^\alpha (1+y)^\beta \left(\frac{(1-y^2)}{dy^2} + [2\beta(1-y) - (2\alpha+1)(1+y)] \frac{d}{dy} + \frac{1}{(1+y)} (2\beta(\beta-1) - 2u_R) - \frac{1}{(1-y)} (2\alpha^2 + 2\varepsilon) - (\alpha^2 + \beta^2) - (\varepsilon + u_0) - u_1 y \right) P_n^{(\mu, \nu)}(y) = 0 \quad (10)$$

The second order differential equation of the Jacobi Polynomial $P_n^{(\mu, \nu)}(y)$ is

$$(1-y^2) \frac{d^2}{dy^2} P_n^{(\mu, \nu)}(y) = \left\{ \begin{aligned} &[(\mu + \nu + 2)y + \mu - \nu] \frac{d}{dy} \\ &-n(n + \mu + \nu + 1) \end{aligned} \right\} \frac{d}{dy} \quad (11)$$




Fig 1. Plot of the potential function given by (1) with $\lambda = 1$. (a) is obtained by vary $u_r = 1, \dots, 11$ (in step of 2) while keeping $u_1 = -5$ and $u_0 = -2$. (b) is obtained by $u_1 = -3$ and $u_0 = -2$ while $u_r = 1, \dots, 11$ (in steps of 2). It is observed that further variations in u_1 and u_0 will result in an upper shift of the potential function in positive xy plane.

Therefore equation (10) becomes

$$\begin{aligned} \frac{2}{\lambda^2} J|\phi_n\rangle &= -A_n (1-y)^\alpha (1+y)^\beta \left(\left[\mu - v + y(\mu + v + 2) \right] \frac{d}{dy} + [2\beta(1-y) - (2\alpha + 1)(1+y)] \frac{d}{dy} + \frac{1}{(1+y)} (2\beta(\beta - 1) - 2u_r) \right) P_n^{(\mu, \nu)}(y) \\ &= -A_n (1-y)^{\mu/2} (1+y)^{v+1/2} \left(\frac{1}{2(1+y)} (v^2 - 1 - 4u_r) + \frac{1}{2(1-y)} (\mu^2 + 4\varepsilon) - (\varepsilon + u_0) - u_1 y - n \left(n + \frac{\mu + v + 1}{2} \right) \right) P_n^{(\mu, \nu)}(y) \end{aligned} \quad (12)$$

Since the matrix representation of the wave operator is required to be tridiagonal and symmetric, in line with the recursion relation of the Jacobi polynomial and its orthogonality; we eliminate the two non-linear terms in (12). Hence the basis parameters must be chosen as follows:

$$v^2 = 1 + 4u_r \text{ and } \mu^2 = -4\varepsilon \quad (13)$$

It is explicit here that the solution of this problem will give negative energy $\mu^2 = -8E/\lambda^2$ and the potential parameter V_r ($u_r \geq -1/4$) should be greater than or equal to zero. Now equation (12) becomes

$$\frac{2}{\lambda^2} J|\phi_n\rangle = -A_n (1-y)^{\mu/2} (1+y)^{v+1/2} \left(-(\varepsilon + u_0) - u_1 y - \left(n + \frac{\mu + v + 1}{2} \right)^2 \right) P_n^{(\mu, \nu)}(y) \quad (14)$$

Using the three term recursion relation of the Jacobi polynomial and their orthogonality property,

$$A_n A_m \int_{-1}^{+1} (1-y)^\mu (1+y)^\nu P_n^{(\mu, \nu)}(y) P_m^{(\mu, \nu)}(y) dy = \delta_{nm} \text{ in (14) after operating } \langle \phi_m | \text{ on the L.H.S of it; we have the tridiagonal and symmetric representation of the wave operator as}$$

$$\frac{2}{\lambda^2} J_{nm} = \left[\left(n + \frac{\mu + \nu + 1}{2} \right)^2 + (\varepsilon + u_0) + u_1 C_n \right] \delta_{nm} + u_1 (D_{n-1} \delta_{n,m+1} + D_n \delta_{n,m-1}) \quad (15)$$

$$\text{where } C_n = \frac{\nu^2 - \mu^2}{(2n + \mu + \nu)(2n + \mu + \nu + 2)} \text{ and } D_n = \frac{2}{2n + \mu + \nu + 2} \sqrt{\frac{(n+1)(n+\mu+1)(n+\nu+1)(n+\mu+\nu+1)}{(2n+\mu+\nu+1)(2n+\mu+\nu+3)}}$$

Hence, we can now write the matrix wave equation as $\langle \phi_n | J | \psi \rangle = \sum_m \langle \phi_n | J | \phi_m \rangle f_m = \sum_m J_{nm} f_m = 0$ which give the three-term recursion relation for the expansion coefficient of the wave function as

$$-(\varepsilon + u_0) f_n = \left[\left(n + \frac{\mu + \nu + 1}{2} \right)^2 + u_1 C_n \right] f_n + u_1 (D_{n-1} f_{n-1} + D_n f_{n+1}) \quad (16)$$

writing $G_n(E) = G_0(E) f_0(\varepsilon)$, will make $f_0 = 1$ and $f_1 = \frac{(z - a_0) f_0}{b_0}$ where $z = -(\varepsilon + u_0)$,

$a_0 = \left(\frac{\mu + \nu + 1}{2} \right)^2 + u_1 C_0$, and $b_0 = u_1 D_0$. This relation is valid for $n = 1, 2, 3, \dots$

This is a new polynomial that we discovered recently and is not found in any mathematics literature. Hence, its analytic properties, that is, weight function, generating function, orthogonality, zero, etc. are yet to be known. Therefore we resolved to a numerical techniques to calculate the energy spectrum of the potential given in (1) for a given set of parameters.

2. Energy spectrum and wavefunction

To calculate the energy spectrum, we obtain first the Hamiltonian matrix from the wave operator matrix (15) as $H = J|_{E=0}$. Then, the energy spectrum is calculated from the wave equation $H|\psi\rangle = E|\psi\rangle$ as the generalized eigenvalues $\{E\}$ of the matrix equation $\sum_m H_{n,m} f_m = E \sum_m \Omega_{n,m} f_m$; where Ω_{nm} is the overlap basis element given as

$$\begin{aligned} \Omega_{n,m} &= \langle \phi_n | \phi_m \rangle = A_n A_m \int_{-1}^1 (1-y)^\mu (1+y)^\nu P_n^{(\mu,\nu)}(y) P_m^{(\mu,\nu)}(y) \times \left[(1-y)^{2\alpha-\mu-a} (1+y)^{2\beta-\nu-b} \right] dy \\ &\equiv \langle n | (1-y)^{2\alpha-\mu-a} (1+y)^{2\beta-\nu-b} | m \rangle \end{aligned} \quad (17)$$

Using the conditions for the parameters of the basis element $2\alpha = \mu$ and $2\beta = \nu + 1$ with $a = 1$ and $b = 0$; the overlap basis element becomes

$$\Omega_{nm} = \langle n | \frac{(1+y)}{(1-y)} | m \rangle = \frac{(1+C_n) \delta_{nm} + D_{n-1} \delta_{n,m+1} + D_n \delta_{n,m-1}}{(1-C_n) \delta_{nm} - D_{n-1} \delta_{n,m+1} - D_n \delta_{n,m-1}} \quad (18)$$

Table 1 is a list of the energy spectrum for a given set of values of the potential parameters and for basis size of $N = 20$. We show only significant decimal digits that do not change with any substantial increase in the basis size (e.g. from size 10 to 50). Also, variation (increase) in N ; we observed a rapid convergence of these values with the size of the basis size

In Figure 2, we plot the bound state wavefunction corresponding to the physical configuration and energy spectrum of Table 1. We calculate the m^{th} bound state using the sum $\psi(E_m, x) \sim \sum_{n=0}^{N-1} P_n(\varepsilon_m) \phi_n(x)$, for some eigenvalues, where N is some large enough integer. An energy resonance (scattering phase shift) phenomenon is observed with this potential function, which will be observed when plotting the wavefunction using some of the eigenvalues in table 1. However, this can be further study using **complex rotation method** [15-17]

3. Conclusion

In this paper, in effort to enlarge the class of solvable potentials, we solve a new potential function that has not been solved in the physics literature to the best of our knowledge. This potential is of greater importance and we will encourage readers to put this into applications. The problem was formulated using the **Tridiagonal Representation Approach**. This method (algebraic in nature) is more encompassing than the conventional methods in solving the Schrödinger equation and allows for more realization of unknown potential functions which are yet to be associated with any physical systems. It has been shown that this approach has contributed greatly to the class of solvable quantum mechanical systems [8]

Table 1: The finite energy spectrum for the potential function (1) with $u_0 = -6$, $u_1 = 10$, and $u_R = 2.5$. It was obtained by diagonalizing the Hamiltonian matrix for different basis size.

n	$\mathcal{E}_n$	n	$\mathcal{E}_n$
0	4126.9891447498	10	32.4394977694
1	1542.8903686294	11	22.4398560240
2	787.2186135745	12	15.2463797234
3	462.6214937613	13	10.1007245429
4	294.0660278716	14	6.4695140302
5	195.9554935304	15	3.9657920559
6	134.3972745542	16	2.2930653158
7	93.7323498172	17	0.0664830132
8	65.8910440926	18	0.4757637553
9	46.3583338365	19	1.1960489428




Fig.2. The graph of the wavefunction as we move down (however there coexist bound states and resonance) the table 1 with physical parameters: $\lambda = 1$, $u_0 = -6$, $u_1 = 10$, and $u_R = 2.5$.

Acknowledgement

The author highly appreciates the support of the Saudi Centre for Theoretical Physics during the progress of this work. Specifically, this paper is dedicated to **Prof. A.D. Alhaidari** for his fatherly role and support in impacting knowledge. Allah in His infinite mercy will continue to bless him more.

Appendix: The Jacobi Polynomials and Manipulations of the Basis Elements

Below are the useful properties of the Jacobi polynomial and the effects of the relevant differential operators on the basis element (5). These are identities are available in textbooks and monographs on orthogonal polynomials [10]

The Jacobi polynomials $P_n^{(\mu,\nu)}(y)$, where $y \in [-1, +1]$ and $\mu > -1$, $\nu > -1$:

$$yP_n^{(\mu,\nu)}(y) = \frac{\nu^2 - \mu^2}{(2n + \mu + \nu)(2n + \mu + \nu + 2)} P_n^{(\mu,\nu)}(y) + \frac{2(n + \mu)(n + \nu)}{(2n + \mu + \nu)(2n + \mu + \nu + 1)} P_{n-1}^{(\mu,\nu)}(y) \\ + \frac{2(n + 1)(n + \mu + \nu + 1)}{(2n + \mu + \nu + 1)(2n + \mu + \nu + 2)} P_{n+1}^{(\mu,\nu)}(y) \quad (\text{A1})$$

$$P_n^{(\mu,\nu)}(y) = \frac{\Gamma(n + \mu + 1)}{\Gamma(n + 1)\Gamma(\mu + 1)} {}_2F_1\left(-n, n + \mu + \nu + 1; \mu + 1; \frac{1-y}{2}\right) = (-1)^n P_n^{(\mu,\nu)}(-y) \quad (\text{A2})$$

$$\left\{ (1-y^2) \frac{d^2}{dy^2} - [(\mu + \nu + 2)y + \mu - \nu] \frac{d}{dy} + n(n + \mu + \nu + 1) \right\} P_n^{(\mu,\nu)}(y) = 0 \quad (\text{A3})$$

$$(1-y^2) \frac{d}{dy} P_n^{(\mu,\nu)}(y) = -n \left(y + \frac{\nu - \mu}{2n + \mu + \nu} \right) P_n^{(\mu,\nu)}(y) + 2 \frac{(n + \mu)(n + \nu)}{2n + \mu + \nu} P_{n-1}^{(\mu,\nu)}(y) \quad (\text{A4})$$

$$\int_{-1}^{+1} (1-y)^\mu (1+y)^\nu P_n^{(\mu,\nu)}(y) P_m^{(\mu,\nu)}(y) dy = \frac{2^{\mu+\nu+1}}{2n + \mu + \nu + 1} \frac{\Gamma(n + \mu + 1)\Gamma(n + \nu + 1)}{\Gamma(n + 1)\Gamma(n + \mu + \nu + 1)} \delta_{nm} \quad (\text{A5})$$

Using the above, with some algebraic manipulations, the effect of differential operators on the basis elements give:

$$(1-y^2) \frac{d^2 \phi_n}{dy^2} = \left[-2\alpha\beta - n(n + \mu + \nu + 1) - n \left(y + \frac{\nu - \mu}{2n + \mu + \nu} \right) \left(\frac{2\alpha - \nu - 1}{1+y} + \frac{1 + \mu - 2\beta}{1-y} \right) + \beta(\beta - 1) \frac{1+y}{1-y} + \alpha(\alpha - 1) \frac{1-y}{1+y} \right] \phi_n \\ + 2 \frac{(n + \mu)(n + \nu)}{2n + \mu + \nu} \left(\frac{2\alpha - \nu - 1}{1+y} + \frac{1 + \mu - 2\beta}{1-y} \right) \frac{A_n}{A_{n-1}} \phi_{n-1} \quad (\text{A6})$$

$$\frac{d\phi_n}{dy} = \left[\frac{\alpha}{1+y} - \frac{\beta}{1-y} - \frac{n}{1-y^2} \left(y + \frac{\nu - \mu}{2n + \mu + \nu} \right) \right] \phi_n + \frac{2}{1-y^2} \frac{(n + \mu)(n + \nu)}{2n + \mu + \nu} \frac{A_n}{A_{n-1}} \phi_{n-1} \quad (\text{A7})$$

The matrix elements of an integrable function $F(y)$ is defined as

$$\langle n | F(y) | m \rangle = A_n A_m \int_{-1}^{+1} (1-y)^\mu (1+y)^\nu F(y) P_n^{(\mu,\nu)}(y) P_m^{(\mu,\nu)}(y) dy \quad (\text{A8})$$

then we obtain

$$\langle n | y | m \rangle = \frac{\nu^2 - \mu^2}{(2n + \mu + \nu)(2n + \mu + \nu + 2)} \delta_{nm} + \frac{2}{(2n + \mu + \nu)} \sqrt{\frac{n(n + \mu)(n + \nu)(n + \mu + \nu)}{(2n + \mu + \nu - 1)(2n + \mu + \nu + 1)}} \delta_{n,m+1} \\ + \frac{2}{(2n + \mu + \nu + 2)} \sqrt{\frac{(n + 1)(n + \mu + 1)(n + \nu + 1)(n + \mu + \nu + 1)}{(2n + \mu + \nu + 1)(2n + \mu + \nu + 3)}} \delta_{n,m-1} \quad (\text{A9})$$

as a special case

Reference

- [1] A.D. Alhaidari and T.J Taiwo, "Four – parameter potential box with inverse square singular boundaries" .Eur. Phys. J. Plus 133 (2018) 115 <https://arxiv.org/abs/1708.05767>.
- [2] A.D. Alhaidari, "An extended class of L^2 - series solution of the wave equation," Ann. Phys.,**317**, 152 2005.
- [3] A.D. Alhaidari, " Analytic solution of the wave equation for an eletron in the field of a molecule with an electric dipole moment, " Ann. Phys., **323**, 1709 2008.
- [4] A.D. Alhaidari, "Scattering and bound states for a class of non –central potentials", J.Phys. A: Math. Theor., **38**. 3409, 2005.
- [5] A.D. Alhaidari, "A class of singular logarithmic potentials in a box with different skin thickness and wall interactions," Phys. Scr., **82**, 065008, 2010.
- [6] A.D. Alhaidari, "Exact solutions of Dirac and Schrodinger equations for a large class of power- law potentials at zero energy," Int. J. Mod. Phys.. A **17**, 4551, 2002.
- [7] P.C. Ojihe, "SO(2,1) Lie algebra, the Jacobi matrix and scattering states of the Morse oscillator," J. Phys. A: Math. Gen., **21**, 875, 1988.
- [8] A.D. Alhaidari, " On the asymptotic solutions of the scattering problem", J. Phys. A: Math. Theor., **41** 175201, 2008.
- [9] A.D. Alhaidari, Solution of the nonrelativistic wave equation using the tridiagonal representation approach, J. Math. Phys., **58**, 072104, 2017.
- [10] R.W. Haymaker and L. Schlessinger, The Pade Approximation in Theoretical Physics, edited by G.A. Baker and J.L Gammel (Academic Press, New York, 1970)
- [11] R. Koekoek and R. Swarttouw, The Askey –Scheme of Hypergeometric orthogonal polynomials and its q analogues, Reports of the Faculty of Technical Mathematics and Informatics, Number 98 -17 (Delft University of Technology, Delft, 1998)
- [12] A.D. Alhaidari and H. Bahlouli, Extending the class of solvable potentials I. The infinite potential well with a sinusoidal bottom, J. Math. Phys.,**49**, 082102, 2008.
- [13] J.S. Geronimo and K. M. Case, Scattering theory and polynomials orthogonal on the real line,Tans. Am. Math., Soc**258**, 467 – 494, 1980.
- [14] F.W.J. Oliver, Asymptotic and Special Functions, (Academic Press, New York, 1974)
- [15] J.S. Geronimo, A relation between the coefficient in the recurrence formula and the spectral Function for the orthogonal polynomials, Tran. Am. Math. Soc., **260**, 65 -82, 1980.
- [16] H .Kamada, Y. Koike, and W. Glockle, "Complex energy method for scattering processes", Prog. Theor. Phys**109**, 869, 2003.
- [17] N. Moiseyev, "Quantum theory of resonances: calculating energies, widths and cross sections by complexscaling ", Phys Rep.,**302**, 212, 1988.
- [18] A. Bottino, A. Longoni, and A. Regge, "Potential scattering for complex energy and angular momentumII NuovoCimento (1995-1965),**23**, 954,1962.



Tunde Joseph Taiwo received his PhD degree in applied mathematics at the University of Benin in Nigeria. He is a member of Saudi Centre for Theoretical Physics (SCTP), International Association of Mathematical Physics (IAMP), Mathematical Association of Nigeria (MAN), and Science Teacher Association of Nigeria (STAN)

His research interests are in the areas of applied mathematics and mathematical physics including: Theoretical quantum mechanics, special functions and orthogonal polynomials, non-Hermitian quantum mechanics and super - symmetric quantum mechanics, mathematical physics - Integrability and J- matrix method. He has published research articles in reputed international journals of theoretical/mathematical physics and mathematical sciences.