

Preliminary Phytochemical and Anti-Bacterial Sensitivity Test of Ethanolic Stem Bark Extract of *Acanthus ilicifolius* (L.)

Madhuri Mondal, Muhammad Torequl Islam*, Shanita Zaman Smrity, and Razina Rouf

Department of Pharmacy, Life Science Faculty, Bangabandhu Sheikh Mujibur Rahman Science and Technology University, Gopalganj-8100, Bangladesh

Received: 8 Sep, 2020, Revised: 6 Nov. 2020, Accepted: 7 Dec. 2020.

Published online: 1 Jan. 2021.

Abstract: This study aimed at the investigation of phytochemical and anti-bacterial activity of *Acanthus ilicifolius* (L.). For this, the qualitative identification of phytochemicals was done in the crude ethanolic stem bark extract of the plant. The anti-bacterial sensitivity test was performed by using disc diffusion and serial dilution methods against *Escherichia coli*, *Bacillus megaterium*, *Bacillus subtilis*, *Staphylococcus entericus* and *Staphylococcus aureus*. Results suggest that the crude extract revealed the presence of alkaloids, flavonoids, terpenoids, phenols, glycosides, steroids, tannins, carbohydrates, fixed oils and fats. The extract at 500 µg/disc produced 14 and 12 mm zones of inhibition against *B. subtilis* and *B. megaterium*, respectively. The minimum inhibitory concentrations calculated for *B. subtilis* and *B. megaterium* was 46.875 µg/mL. In conclusion, the ethanolic stem bark extract of *A. ilicifolius* showed an inhibitory effect against the test bacterial strains, suggesting the plant may be a good source of anti-bacterial agents.

Keywords: *Acanthus ilicifolius*, phytochemicals, anti-bacterial effect.

1 Introduction

Acanthus ilicifolius (L.) (Sea holly) (Family: Acanthaceae) is a perennial herbaceous mangrove plant, popularly recognized as “Holy leaved acanthus” or “Kayalchulli”. Traditionally *A. ilicifolius* is known as Sahachara and used in Indian Ayurveda. *A. ilicifolius* is popularly used for its wound healing ability by the coastal inhabitants of West Bengal. The root is expectorant, and is used in coughs and asthma. The plant normally lives in areas of modest salinity, forming bush around mangrove palms. From very ancient time, the plant is applied in traditional systems of medicine, including traditional Indian medicine or Ayurveda and traditional Chinese medicine [1,2] in the intend of medicating various diseases. Different parts of the plant have been used to treat asthma, diabetes, dyspepsia, leprosy, hepatitis, paralysis, snake bite, rheumatoid arthritis and are used as diuretics [3-5].

Phytochemical studies with the plant revealed the presence of lignans [6] and megastigmane glycosides [7]. Scientific reports suggest that the plant has antioxidant, anti-inflammatory, anti-leishmanial, osteoblastic, hepatoprotective, anticancer, antiulcer, analgesic and antimicrobial activities [1,4,5,8-10].

Drug resistance in human pathogenic microorganisms is a continuous process, which leads to search new antimicrobial drugs. These circumstances imposed scientists to search for new anti-microbial substances from various sources, including medicinal plants [1]. This study aimed to investigate anti-bacterial activity of ethanolic bark extract of *A. ilicifolius* against a number of human pathogenic bacterial strains along with the preliminary phytochemical screening.

2 Experimental Section

2.1 Collection and identification of plant material

The fresh stem bark of the *A. ilicifolius* was collected from the Adampur reserve forest, Kamalgonj, Moulvibazar, Bangladesh in the month of July, 2019 and was identified by the taxonomist at the Department of Botany, Jahangirnagar University, Bangladesh. A voucher specimen (Accession number-DACB: 48258) was deposited in the Bangladesh National Herbarium, Dhaka (Mirpur), Bangladesh.

* Corresponding author E-mail: dmt.islam@bsmrstu.edu.bd

2.2 Drying and grinding

The collected stem bark was separated from undesirable materials, washed with running tap water and shade-dried for one week. Then the plant materials were ground into a coarse powder with the help of a suitable grinder (Capacitor start motor, Wuhu motor factory, China). The powder was stored in an airtight container and kept in a cool, dark and dry place until the test commenced.

2.3 Cold extraction (maceration)

The plant was extracted by the cold extraction or maceration method. About 250 g of powder was soaked in 900 mL of 99.9% ethanol in an amber color glass container for 9 days. Then the extract was concentrated by solvent evaporation at room temperature and dried to a solid in an oven (temperature not exceeding 50 °C).

2.4 Phytochemical screening

In preliminary phytochemical screening test for secondary metabolites such as alkaloids [11,12], flavonoids [12,13], triterpenoids [14,15], tannins [16,17], carbohydrates [12,18], glycosides [13,19,20], phenolic compounds, amino acids and proteins [11], steroids [14], fixed oil and fats [18] were done.

2.5 Anti-bacterial sensitivity test

The agar diffusion assay was done according to Bonev *et al.* [21], while the minimum inhibitory concentration test was done according to Andrews [22].

3 Results and Discussion

35.36 mg greenish color type of ethanolic crude extract was found. The yield value of the plant extract was 0.14. Phytochemicals can protect against disease and are the good sources of modern medicine [23]. In this study, the phytochemical screening report suggests that the crude extract possesses alkaloids, flavonoids, terpenoids, phenols, glycosides, steroids, tannins, carbohydrates, fixed oils and fats (Table 1). Generally, the secondary metabolites of medicinal plants are the promising source of many modern drugs [24]. Flavonoids and phenols are responsible for the antioxidant and hepatoprotective effects [23]. Babu *et al.* [4] also previously reported that *A. ilicifolius* has antioxidant and hepatoprotective activity.

Table 1. Phytochemical relevance of ethanolic stem bark extract of *A. ilicifolius*

Phytochemical groups	Tests	Relevance
Alkaloids	Dragendroff's test	-

	Mayer's test	+
	Wagner's test	+
	Hager's test	+
Flavonoids	Alkaline reagent	++
	Shinoda's test	++
	Lead acetate test	+++
Terpenoids	Salkowski test	++
Tannins	Ferric Chloride test	+++
	Lead sub acetate test	++
Carbohydrates	Molisch test	++
	Benedict's test	+++
	Fehling's test	+++
Glycosides	Legal test	++
	Keller-killiani	+++
	Borntrager's test	++
Phenols	Ferric chloride test	+++
Amino acids and proteins	Biuret test	-
	Million test	-
Steroids	Libermann-Burchard's test	++
Fixed oils and fats	Spot test	+++
Intensity: + = Trace; ++ = Moderate; +++ = Strong; - = Absence		

In a study, the alcoholic and chloroform extracts of the leaves of the plant were found to exhibit strong inhibitory activity against *B. subtilis*, *S. aureus*, *Candida albicans*, *Aspergillus fumigatus* and *Aspergillus niger*, while moderate inhibitory action against *Pseudomonas aeruginosa* and *Proteus vulgaris* [25]. In another study, the chloroform extract showed maximum activity against the bacterial pathogens methicillin-resistant *S. aureus*, *Streptococcus pyogenes*, *Pseudomonas aeruginosa*, *C. albicans* and *Trichophyton rubrum*. Methanol and acetone extracts showed maximum activity against *S. epidermis* and *Lactobacillus plantarum*, respectively [26]. The plant is also evident to act against *Aspergillus fumigatus* [27]. In this study, the ethanolic stem bark extract produced highest zone of inhibition (14 mm) against *B. subtilis*, then followed by *B. megaterium*, *S. aureus*, *E. coli* and *S. entericus*, respectively. The crude extract of the plant showed lowest the MIC value (46.875 µg/mL) against *B. subtilis* and *B. megaterium*. It showed MIC value 187.5 µg/mL against *S. aureus*. Against other two strains, *E. coli* and *S. entericus*, the extract showed 750 µg/mL MIC value (Table 2 & Figure 1).

Table 2. Anti-bacterial activity of ethanolic stem bark extract of *A. ilicifolius*

Bacteria	ZI (mm)		MIC value
	Azithromycin (30 µg/disc)	Crude extract (500)	Crude extract (µg/mL)

		$\mu\text{g/disc}$	
<i>Bacillus megaterium</i>	33	12	46.875
<i>Bacillus subtilis</i>	30	14	46.875
<i>Escherichia coli</i>	13	7	750
<i>Staphylococcus aureus</i>	15	9	187.5
<i>Staphylococcus entericus</i>	16	7	750
ZI: Zone of inhibition; MIC: Minimum inhibitory concentration			


*E. coli**S. aureus**B. megaterium*

Figure 1. Zone of inhibition of the plant extract against some test bacterial strains

4 Conclusions

The ethanolic crude extract of *A. ilicifolius* possesses many important secondary metabolites, including alkaloids, flavonoids, terpenoids, phenols, glycosides, steroids, and tannins. The extract showed moderate anti-bacterial activity against *B. subtilis* and *B. megaterium*. Scientific reports also suggest that various parts of the plant also possess broad-spectrum anti-microbial effect. Further research is required to isolate the active compounds responsible for its anti-microbial effects.

Acknowledgement

We are owed to the Department of Pharmacy, BSMRSTU, Gopalganj (Bangladesh) for providing laboratory facilities to conduct this study. We are also thankful to the Pharmacy Discipline for providing the test bacterial strains for this study.

Conflict of interest

None declared.

References

[1] R. Wöstmann, G. Liebezeit, Chemical composition of the mangrove holly *Acanthus ilicifolius* (Acanthaceae)—review and additional data, *Senckenbergianamaritima*, **38(1)**, 31, 2008.

- [2] L. Liu, H. Fan, P. Qi, Y. Mei, L. Zhou, L. Cai, X. Lin, J. Lin, Synthesis and hepatoprotective properties of *Acanthus ilicifolius* alkaloid A and its derivatives, *Exp Therap Med.*, **6(3)**, 796-802, 2013.
- [3] W. M. Bandaranayake, Traditional and medicinal uses of mangroves, *Mangroves Salt Marshes*, **2(3)**, 133-148, 1998.
- [4] B. H. Babu, B. S. Shylesh, J. Padikkala, Antioxidant and hepatoprotective effect of *Acanthus ilicifolius*, *Fitoterapia*, **72(3)**, 272-277, 2001.
- [5] A. Simlai, A. Roy, Biological activities and chemical constituents of some mangrove species from Sundarban estuary: An overview, *Pharmacogn Rev.*, **7(14)**, 170, 2013.
- [6] T. Kanchanapoom, M. S. Kamel, R. Kasai, K. Yamasaki, C. Picheansoonthon, Y. Hiraga, Lignan glucosides from *Acanthus ilicifolius*, *Phytochemistry*, **56(4)**, 369-372, 2001.
- [7] J. Wu, S. Zhang, Q. Xiao, Q. Li, J. Huang, L. Long, L. Huang, Megastigmane and flavone glycosides from *Acanthus ilicifolius*, *Die Pharmazie – Int J Pharm Sci.*, **58(5)**, 363-364, 2003.
- [8] N. V. Agshikar, V. R. Naik, G. J. Abraham, C. V. Reddy, S. W. Naqvo, P. K. Mittal, Analgesic anti-inflammatory activity of *Acanthus ilicifolius* Linn., *Indian J Exp Biol.*, **17(11)**, 1257-1258, 1979.
- [9] A. Kapil, S. Sharma, S. Wahidulla, Leishmanicidal activity of 2-benzoxazolinone from *Acanthus ilicifolius* in vitro, *Planta Med.*, **60(02)**, 187-188, 1994.
- [10] B. H. Babu, B. S. Shylesh, J. Padikkala, Tumour reducing and anticarcinogenic activity of *Acanthus ilicifolius* in mice, *J Ethnopharmacol.*, **79(1)**, 27-33, 2002.
- [11] A. Kumar, K. K. Jha, D. Kumar, A. Agrawal, A. Gupta, Preliminary phytochemical analysis of leaf and bark (mixture) extract of *Ficus infectoria* plant, *Pharm Innov.*, **1(5, Part A)**, 71, 2012.
- [12] C. S. Vimal kumar, V. B. Hosagaudar, S. R. Suja, V. Vilash, N. M. Krishnakumar, P. G. Latha, Comparative preliminary phytochemical analysis of ethanolic extracts of leaves of *Olea dioica* Roxb., infected with the rust fungus *Zaghouaniaoleae* (E.J. Butler) Cummins and non-infected plants, *J Pharmacogn Phytochem.*, **3(4)**, 69-72, 2014.
- [13] R. N. S. Yadav, M. Agarwala, Phytochemical analysis of some medicinal plants, *J Phylol.*, **3(12)**, 10-14, 2011.
- [14] C. Kodangala, S. Saha, P. Kodangala, Phytochemical studies of aerial parts of the plant *Leucas lavandulaefolia*. Scholars Research Library, *Der Pharm Chem.*, **2(5)**, 434-437, 2010.
- [15] R. S. Kumar, C. Venkateshwar, G. Samuel, S. Gangadhar Rao, Phytochemical Screening of some compounds from plant leaf extracts of *Holoptelea integrifolia* (Planch.) and *Celestrusemarginata* (Grah.) used by Gondu tribes at Adilabad District, Andhrapradesh, India, *Int J Eng Sci Invent.*, **2(8)**, 65-70, 2013.
- [16] S. F. Zohra, B. Meriem, S. Samira, M. S. Alsayadi Muneer, Phytochemical Screening and identification of some compounds from Mallow, *J Nat Prod Plant Resour.*,

2(4), 512-516, 2012.

- [17] V. Ramamurthy, M. Sathiyadevi, Preliminary Phytochemical Screening of Methanol Extract of *Indigo feratrita* Linn., *J Mol Histol Med Physiol.*, **2**, 112, 2017.
- [18] K. L. Singh, G. C. Bag, Phytochemical Analysis And Determination Of Total Phenolics Content In Water Extracts Of Three Species Of *Hedychium*, *Int J Pharm Tech Res CODEN (USA): IJPRIF*, **5(4)**, 1516-1521, 2013.
- [19] G. George, Preliminary Phytochemical Screening of Whole Plant Extracts of *Peperomia pellucida* (Linn.) HBK (Piperaceae) and *Marsilea quadrifolia* Linn. (Marsileaceae), *Int J Pharmacogn Phytochem Res.*, **5(3)**, 200-214, 2013.
- [20] Y. Rufai, Y. Isah, M. S. Isyaka, Comparative Phyto-Constituents Analysis from the Root Bark and Root Core Extractives of *Cassia ferruginea* (Schrad D. C) Plant, *Sch J Agric Vet Sci.*, **3(4)**, 275-283, 2016.
- [21] B. Bonev, J. Hooper, J. Parisot, Principles of assessing bacterial susceptibility to antibiotics using the agar diffusion method, *J Antimicrob Chemother.*, **61(6)**, 1295-301, 2008.
- [22] J. M. Andrews, Determination of minimum inhibitory concentrations, *J Antimicrob Chemother.*, **48(suppl 1)**, 5-16, 2001.
- [23] J. Lee, D. -G. Jo, D. Park, H. Y. Chung, M. P. Mattson, Adaptive Cellular Stress Pathways as Therapeutic Targets of Dietary Phytochemicals: Focus on the Nervous System, *Pharmacol Rev.*, **66(3)**, 815-868, 2014.
- [24] J. Nankaya, N. Gichuki, C. Lukhoba, H. Balslev, Medicinal Plants of the Maasai of Kenya: A Review, *Plants (Basel)*, **9(1)**, 44, 2020.
- [25] S. Bose, A. Bose, Antimicrobial Activity of *Acanthus ilicifolius* (L.), *Indian J Pharm Sci.*, **70(6)**, 821-823, 2008.
- [26] C. Govindasamy, M. Arulpriya, Antimicrobial activity of *Acanthus ilicifolius*: Skin infection pathogens, *Asian Pac J Trop Dis.*, **3(3)**, 180-183, 2003.
- [27] P. S. Kalaskar, V. V. Karande, A. S. Bannaliker, M. M. Gatne, Antifungal Activity of Leaves of Mangroves Plant *Acanthus licifolius* Against *Aspergillus fumigatus*, *Indian J Pharm Sci.*, **74(6)**, 575-579, 2012.