

International Journal of Advanced Biomedicine

A Dynamic Model for Predicting Immune System Interactions with Infectious Diseases

E. L. Pankratov^{1,*}

¹ Department of Applied Mechanics, Physics and Higher Mathematics of Nizhny Novgorod State Agrotechnical University, 97 Gagarin avenue, Nizhny Novgorod, 603950, Russia.

Received: 12 Dec. 2024, Revised: 7 Feb. 2025, Accepted: 29 Mar. 2025. Published online: 1 May 2025.

Abstract: In this paper we consider a model of interaction of the human immune system with infectious diseases with account changes in its conditions. We also consider an analytical approach for analyzing of the above interaction. It has been also considered possibility of acceleration and deceleration of the above interaction.

Keywords: interaction of the human immune system with infectious diseases; changing of conditions of interaction; prognosis of interaction; analytical approach for analysis.

1 Introduction

Currently, the development of immunology is proceeding so rapidly that its concepts are changing before our eyes, involving new facts and hypotheses in the arsenal of ideas about immune processes that correct various elements of the theory [1-5]. Updating information about immune processes leads to the need to correction of the considered treatment program. An actual issue in the correction of the treatment program is the prognosis of changes in the immune system during its control. In this paper we consider a model of the interaction of the human immune system with infectious diseases with account changes in its conditions. We also consider an analytical approach for analyzing of the above interaction. It has been also considered possibility of acceleration and deceleration of the above interaction.

2 Method of Solution

1

In this section we consider the following model of interaction of the human immune system with infectious diseases

$$\begin{cases} \frac{dV(t)}{dt} = \beta(t)V(t) - \gamma(t)F(t)V(t) \\ \frac{dF(t)}{dt} = \rho(t)C(t) - \eta(t)\gamma(t)F(t)V(t) - \mu_f(t)F(t) \\ \frac{dC(t)}{dt} = \xi(t)\alpha(t)F(t-\tau)V(t-\tau) - \mu_c(t)[C(t) - C^*] \\ \frac{dm(t)}{dt} = \sigma(t)V(t) - \mu_m(t)m(t) \end{cases}$$
(1)

where V(t) is the concentration of proliferating antigens; C (t) is the concentration of antigen-specific lymphocytes (carriers and producers of antibodies) in lymphoid tissue; F(t) is the concentration of antibodies (molecules of an immune nature, immunoglobulins, receptors of immunocompetent cells, etc. - neutralizing antigens) in the blood; m(t) is the the proportion of cells destroyed by the antigen in the affected part of the target organ, affecting the weakening of the body's vital functions during the course of the disease, associated with a decrease in the activity of organs that provide the supply of immunological material: leukocytes, lymphocytes, antibodies, etc., necessary to combat multiplying antigens; β (t) is the rate of reproduction of antigens; $\gamma(t)$ is the coefficient that takes into account the probability of encountering viruses with antibodies and the strength of their interaction; α (t) is the immune system stimulation factor; ρ (t) is the the rate of production of antibodies by a single plasma cell; μ_c (t) and μ_f (t) are the values inverse to the life span of plasma cells and antibodies, respectively; η (t) is the the amount of antibodies needed to neutralize a single virus; $\sigma(t)$ is the organ damage rate; $\mu_m(t)$ is the the rate of recovery of the mass of the affected organ; C(t) is the pre-existing level of immunocompetent cells (plasma cells); τ is the time required for the formation of a cascade of plasma cells; ξ (t) is the a function that takes into account the disruption of the normal functioning of the immune system due to significant damage to the organ. The initial conditions of the desired functions described by the equations of system (1) can be represented in the following form

 $V(0) = V_0, F(0) = F_0, C(0) = C_0, m(0) = m_0.$ (2)

Now we transform differential equations of the system (1) to the following integral form with account conditions (2)



$$\begin{cases} V(t) = \int_{0}^{t} \beta(\chi) V(\chi) d\chi - \int_{0}^{t} \gamma(\chi) F(\chi) V(\chi) d\chi + V_{0} \\ F(t) = \int_{0}^{t} \rho(\chi) C(\chi) d\chi - \int_{0}^{t} \eta(\chi) \gamma(\chi) F(\chi) V(\chi) d\chi - \int_{0}^{t} \mu_{f}(\chi) F(\chi) d\chi + F_{0} \\ C(t) = \int_{0}^{t} \xi(\chi) \alpha(\chi) F(\chi - \tau) V(\chi - \tau) d\chi - \int_{0}^{t} \mu_{C}(\chi) [C(\chi) - C^{*}] d\chi + C_{0} \\ m(t) = \int_{0}^{t} \sigma(\chi) V(\chi) d\chi - \int_{0}^{t} \mu_{m}(\chi) m(\chi) d\chi + m_{0} \end{cases}$$
(3)

Next we calculate solutions of equations of the system (3) by the method of averaging of function corrections [6-8]. In the framework of the approach we replace the required functions in the right sides of equations (3) on their not yet known average values α_1 . The substitution gives a possibility to obtain equations to determine the first-order approximations of the required functions in the following form

$$\begin{cases} V_{1}(t) = \alpha_{1V} \int_{0}^{t} \beta(\chi) d\chi - \alpha_{1V} \alpha_{1F} \int_{0}^{t} \gamma(\chi) d\chi + V_{0} \\ F_{1}(t) = \alpha_{1C} \int_{0}^{t} \rho(\chi) d\chi - \alpha_{1F} \alpha_{1V} \int_{0}^{t} \eta(\chi) \gamma(\chi) d\chi - \alpha_{1F} \int_{0}^{t} \mu_{f}(\chi) d\chi + F_{0} \\ C_{1}(t) = \alpha_{1V} \alpha_{1F} \int_{0}^{t} \xi(\chi) \alpha(\chi) F(\chi - \tau) V(\chi - \tau) d\chi - (\alpha_{1C} - C^{*}) \int_{0}^{t} \mu_{C}(\chi) d\chi + C_{0} \\ m_{1}(t) = \alpha_{1V} \int_{0}^{t} \sigma(\chi) d\chi - \alpha_{1m} \int_{0}^{t} \mu_{m}(\chi) d\chi + m_{0} \end{cases}$$

$$(4)$$

Average values α_1 were determined by the following standard relation [6-8]

$$\alpha_1 = \frac{1}{\Theta} \int_0^{\Theta} G_1(t) dt , \qquad (5)$$

where Θ is the continuance of observation on the considered process. Substitution of relations (4) into relation (5) gives a possibility to obtain relations to calculate the required average values α_1 in the following final form

$$\begin{cases} \Theta \alpha_{1V} = \alpha_{1V} \int_{0}^{\Theta} (\Theta - t) \beta(t) dt - \alpha_{1V} \alpha_{1F} \int_{0}^{\Theta} (\Theta - t) \gamma(t) dt + V_{0} \Theta \\ \Theta \alpha_{1F} = \alpha_{1C} \int_{0}^{\Theta} (\Theta - t) \rho(t) dt - \alpha_{1F} \alpha_{1V} \Theta \int_{0}^{\Theta} (\Theta - t) \eta(t) \gamma(t) dt - \alpha_{1F} \int_{0}^{\Theta} (\Theta - t) \mu_{f}(t) dt + \Theta F_{0} \\ \Theta \alpha_{1C} = \alpha_{1V} \alpha_{1F} \int_{0}^{\Theta} (\Theta - t) \xi(t) \alpha(t) dt - (\alpha_{1C} - C^{*}) \int_{0}^{\Theta} (\Theta - t) \mu_{C}(t) dt + \Theta C_{0} \\ \Theta \alpha_{1m} = \alpha_{1V} \int_{0}^{\Theta} (\Theta - t) \sigma(t) dt - \alpha_{1m} \int_{0}^{\Theta} (\Theta - t) \mu_{m}(t) dt + \Theta m_{0} \end{cases}$$

$$\tag{6}$$

Solution of equations of the system (6) by standard approaches [9] could be presented in the following form

$$\alpha_{1c} = \frac{ch - cf^{2} - 2dg - 2eg}{3cg} + \frac{3c^{2}g(2cdf^{2} - 2cef + d^{2}g + 2deg + e^{2}g - 2cdh - 2ceh) - c^{2}(ch - cf^{2} - 2dg - 2eg)^{2}}{4c^{2}g(c^{6}f^{6} + 9bc^{5}ef^{3}g - 3c^{5}df^{4}g)},$$

$$\alpha_{1V} = \frac{V_{0}\Theta}{\Theta - \int_{0}^{\Theta}(\Theta - t)\beta(t)dt + \alpha_{1F}\int_{0}^{\Theta}(\Theta - t)\gamma(t)dt},$$

$$\alpha_{1F} = \frac{\sqrt{b^{2} + 4\alpha_{1C}\int_{0}^{\Theta}(\Theta - t)\rho(t)dt - 4\Theta F_{0}} - b}{2a},$$

$$(7)$$

$$\alpha_{1m} = \frac{V_{0}\Theta\int_{0}^{\Theta}(\Theta - t)\sigma(t)dt + \Theta m_{0}\left[\Theta - \int_{0}^{\Theta}(\Theta - t)\beta(t)dt + \alpha_{1F}\int_{0}^{\Theta}(\Theta - t)\gamma(t)dt\right]}{\left[\Theta - \int_{0}^{\Theta}(\Theta - t)\beta(t)dt + \alpha_{1F}\int_{0}^{\Theta}(\Theta - t)\gamma(t)dt\right]},$$

$$mhere$$

$$b = \left[\Theta - \int_{0}^{\Theta}(\Theta - t)\beta(t)dt \right] \left[\int_{0}^{\Theta}(\Theta - t)\gamma(t)dt \left[\Theta + \int_{0}^{\Theta}(\Theta - t)\mu_{f}(t)dt\right],$$

$$f = 2a\Theta - - 2a\int_{0}^{\Theta}(\Theta - t)\beta(t)dt - 2ab\int_{0}^{\Theta}(\Theta - t)\gamma(t)dt$$

$$c = \Theta + \int_{0}^{\Theta} \mu_{C}(t) (\Theta - t) dt, \qquad e = V_{0} \Theta \int_{0}^{\Theta} (\Theta - t) \xi(t) \alpha(t) dt,$$
$$h = b^{2} - 4\Theta F_{0},$$

$$g = 4 \int_{0}^{\Theta} (\Theta - t) \rho(t) dt , \quad d = C^* \int_{0}^{\Theta} (\Theta - t) \mu_C(t) dt + \Theta C_0$$

The second-order approximations of the required functions in the framework of the method of averaging of function corrections could be calculated by replacement of the above functions on the following sums: $G(t) \rightarrow \alpha_2 + G_1(t)$ [6-8]. The replacement gives a possibility to obtain the following equations to calculate the considered functions

$$\begin{cases} V_{2}(t) = \int_{0}^{t} \beta(\chi) [\alpha_{2v} + V_{1}(\chi)] d\chi - \int_{0}^{t} \gamma(\chi) [\alpha_{2F} + F_{1}(\chi)] [\alpha_{2v} + V_{1}(\chi)] d\chi + V_{0} \\ F_{2}(t) = \int_{0}^{t} \rho(\chi) [\alpha_{2c} + C_{1}(\chi)] d\chi - \int_{0}^{t} \eta(\chi) \gamma(\chi) [\alpha_{2F} + F_{1}(\chi)] [\alpha_{2v} + V_{1}(\chi)] d\chi - \int_{0}^{t} \mu_{f}(\chi) [\alpha_{2v} + V_{1}(\chi)] d\chi + F_{0} \\ C_{2}(t) = \int_{0}^{t} \xi(\chi) \alpha(\chi) [\alpha_{2F} + F_{1}(\chi - \tau)] [\alpha_{2v} + V_{1}(\chi - \tau)] d\chi + C_{0} - \int_{0}^{t} \mu_{c}(\chi) [\alpha_{2c} + C_{1}(\chi) - C^{*}] d\chi \\ m_{2}(t) = \int_{0}^{t} \sigma(\chi) [\alpha_{2v} + V_{1}(\chi)] d\chi - \int_{0}^{t} \mu_{m}(\chi) [\alpha_{2m} + m_{1}(\chi)] d\chi + m_{0} \end{cases}$$

Average values of the second-order approximations α_2 were determined by the following standard relation [6-8]

(8)

^{© 2025} NSP

Natural Sciences Publishing Cor.

Int. J. Adv. Biomed., 9, No. 2, 7-10 (2025)/ http://www.naturalspublishing.com/Journals.asp

$$\alpha_2 = \frac{1}{\Theta} \int_0^{\Theta} [G_2(t) - G_1(t)] dt$$
(9)

Substitution of relations (8) into relations (9) gives a possibility to obtain the following relations to determine the required average values α_2

$$\begin{vmatrix} \Theta \alpha_{2v} &= \int_{0}^{\Theta} (\Theta - t) \beta(t) [\alpha_{2v} + V_{1}(t)] dt + \Theta V_{0} - \int_{0}^{\Theta} (\Theta - t) \gamma(t) [\alpha_{2F} + F_{1}(t)] [\alpha_{2v} + V_{1}(t)] dt \\ \Theta \alpha_{2F} &= \int_{0}^{\Theta} (\Theta - t) \rho(t) [\alpha_{2C} + C_{1}(t)] dt - \int_{0}^{\Theta} (\Theta - t) \eta(t) \gamma(t) [\alpha_{2F} + F_{1}(t)] [\alpha_{2v} + V_{1}(t)] dt + \Theta F_{0} - \\ \left\{ - \int_{0}^{\Theta} (\Theta - t) \mu_{f}(\chi) [\alpha_{2v} + V_{1}(t)] dt \\ \Theta \alpha_{2C} &= \int_{0}^{\Theta} (\Theta - t) \xi(t) \alpha(t) [\alpha_{2F} + F_{1}(t - t)] [\alpha_{2v} + V_{1}(t - t)] dt - \int_{0}^{\Theta} (\Theta - t) \mu_{c}(t) [\alpha_{2C} + C_{1}(t) - C^{*}] dt + \Theta C_{0} \\ \Theta \alpha_{2m} &= \int_{0}^{\Theta} (\Theta - t) \sigma(t) [\alpha_{2v} + V_{1}(t)] dt - \int_{0}^{\Theta} (\Theta - t) \mu_{m}(t) [\alpha_{2m} + m_{1}(t)] dt + \Theta m_{0} \end{aligned}$$

$$(10)$$

Solution of equations of the system (10) by standard approaches [9] could be presented in the following form

$$\alpha_{2m} = \frac{\bigcap_{0}^{\Theta}(\Theta - t)\sigma(t)[\alpha_{2V} + V_{1}(t)]dt - \bigcap_{0}^{\Theta}(\Theta - t)\mu_{m}(t)m_{1}(t)dt + \Theta m_{0}}{\Theta + \bigcap_{0}^{\Theta}(\Theta - t)\mu_{m}(t)dt}$$

$$\alpha_{2V} = \frac{p - \alpha_{2F} \bigcap_{0}^{\Theta}(\Theta - t)\gamma(t)V_{1}(t)dt}{\left[q + \alpha_{2F} \bigcap_{0}^{\Theta}(\Theta - t)\gamma(t)dt\right]},$$

$$\alpha_{2F} = \frac{\alpha_{2C} - A}{3x} - \frac{\sqrt[3]{2}(D - E\alpha_{2C})}{2A^{2}(A + 2\alpha_{2C})},$$

$$\alpha_{2C} = \frac{1}{2Ep} \left[A^{2} \left\{\bigcap_{0}^{\Theta}(\Theta - t)\mu_{C}(t)[C_{1}(t) - C^{*}]dt - \Theta C_{0}\right\} + 2 \left[A^{2} \left\{\bigcap_{0}^{\Theta}(\Theta - t)\xi(t)\alpha(t)dt\right\} - \left[\Theta + \bigcap_{0}^{\Theta}(\Theta - t)\mu_{C}(t)dt\right]\right]$$

$$\times \left[\bigcap_{0}^{\Theta}(\Theta - t)\xi(t)\alpha(t)F_{1}(t - \tau)dt + \bigcap_{0}^{\Theta}(\Theta - t)\xi(t)\alpha(t)V_{1}(t - \tau)dt\right]^{-1} \left[\Theta + \bigcap_{0}^{\Theta}(\Theta - t)\mu_{C}(t)dt\right]^{-1},$$

$$(11)$$

$$\alpha = \Theta - \bigcap_{0}^{\Theta}(\Theta - t)\beta(t)dt + \bigcap_{0}^{\Theta}(\Theta - t)\beta(t)dt + \bigcap_{0}^{\Theta}(\Theta - t)\gamma(t)E(t)dt$$

where

$$p = \int_{0}^{\Theta} (\Theta - t) \beta(t) V_1(t) dt + \Theta V_0 - \int_{0}^{\Theta} (\Theta - t) \gamma(t) F_1(t) V_1(t) dt$$

$$r = \int_{0}^{\Theta} (\Theta - t) \rho(t) C_{1}(t) dt - \int_{0}^{\Theta} (\Theta - t) \mu_{f}(\chi) V_{1}(t) dt + \Theta F_{0} - \int_{0}^{\Theta} (\Theta - t) \eta(t) \gamma(t) F_{1}(t) V_{1}(t) dt$$
$$u = \int_{0}^{\Theta} (\Theta - t) V_{1}(t) \times$$

$$\begin{split} & \times \beta(t)dt - \int_{0}^{t} (\Theta - t)\gamma(t)F_{1}(t)V_{1}(t)dt + \Theta V_{0}, \\ & s = \Theta + \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)V_{1}(t)dt \\ & , \\ & v = \Theta - \int_{0}^{\theta} (\Theta - t)\beta(t)dt + \int_{0}^{\theta} (\Theta - t)F_{1}(t) \times \\ & \times \gamma(t)dt, \\ & x = \int_{0}^{\theta} (\Theta - t)\gamma(t)dt \left[\int_{0}^{\theta} (\Theta - t)\gamma(t)dt - \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t)dt \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)dt \right] \\ & y = v \int_{0}^{\theta} (\Theta - t)\gamma(t) dt - \\ & -v \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t)dt \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)dt + \left[q + p \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)dt \right] \\ & y = v \int_{0}^{\theta} (\Theta - t)\gamma(t) dt - \\ & -v \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t)dt \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)dt + \left[q + p \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)dt \right] \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t)dt + \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t)V_{1}(t)dt \\ & \int_{0}^{\theta} (\Theta - t)\eta(t)\gamma(t) F_{1}(t)dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & = \int_{0}^{\theta} (\Theta - t)\rho(t)dt \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t)dt \\ & = v \left[q + p \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \right] + q w \int_{0}^{\theta} (\Theta - t)\gamma(t)V_{1}(t) dt \\ & = q \int_{0}^{\theta} (\Theta - t)\rho(t)dt \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t)dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \times \gamma \quad (t) dt + w u \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \times \gamma \quad (t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & + v \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\rho(t) dt \\ & \int_{0}^{\theta} (\Theta - t)\gamma(t) \\ & \int_{0$$

Analysis of the considered function has been done analytically by using the second-order approximation in the framework of method of averaging of function corrections. The approximation is usually enough good approximation for to make qualitative analysis and to obtain some quantitative results. All obtained results have been checked by comparison with results of numerical simulations.

3 Discussion

In this section we analyzed the considered concentrations and proportion of cells destroyed by antigen. Figs. 1, 2 and 3 show typical dependences of the considered concentrations on time for developing disease, fading disease and stabilizing disease at different values of the considered parameters. Changing the values of the model parameters with time gives a possibility to accelerate or 10

slow down the development or stabilization of the disease, as well as the recovery of the body or change the mode of development of the situation.



Fig. 1: Typical dependences of the concentration of proliferating antigens on time for developing disease







Fig. 3: Typical dependences of the concentration of proliferating antigens on time for stabilizing disease

© 2025 NSP Natural Sciences Publishing Cor.

5 Conclusions

We consider a model of interaction of the human immune system with infectious diseases with account changes in its conditions. Also we consider an analytical approach for analyzing of the above interaction. It has been also considered possibility to change speed of the above interaction.

References

- G.A. Bocharov, A.A. Danilov, Yu.V. Vasilevsky, G.I. Marchuk, V.A. Chereshnev, B. Lyudevig, Modeling of the protective layer of the interferon field in lymphoid organs taking into account their structural and functional organization, Reports of Academy of Science of Russia, 439 (3), 413-415, 2011.
- [2] E.N. Severskaya, T.E. Eltyshova, D.S. Karpov, M.P. Korobkina, P.G. Zaikina, Lipid analysis of lacrimal fluid in the diagnosis of meibomian gland dysfunction, Russian medical journal, Issue 1, 9-14, 2022.
- [3] O.A. Shavlovskaya, Evaluation of the dynamics of Ddimer and C-reactive protein levels in patients with osteoarthritis who have had COVID-19, during therapy with parenteral chondroitin sulfate, Russian medical journal, Issue 7, 32-38, 2022.
- [4] D. Dasgupt. Artificial immune systems and their applications (Moscow: FISMATLIT, 2006).
- [5] S.R. Kuznetsov, Mathematical model of immune response, Bulletin of Saint Petersburg State University, Issue 4, 72-87, 2015.
- [6] Yu.D. Sokolov, About the definition of dynamic forces in the mine lifting, Applied Mechanics, 1 (1), 23-35, 1955.
- [7] E.L. Pankratov, E.A. Bulaeva, An approach to decrease dimensions of field-effect transistors without *p-n*junctions, International Journal of Modern Physics B, 28 (27), 1450190-1--1450190-17, 2014.
- [8] E.L. Pankratov, E.A. Bulaeva, On optimization of regimes of epitaxy from gas phase. some analytical approaches to model physical processes in reactors for epitaxy from gas phase during growth films, Reviews in Theoretical Science, 3 (4), 365-398, 2015.
- [9] G. Korn, T. Korn. Handbook of Mathematics for Scientists and Engineers (Moscow: Nauka, 1974).