Applied Mathematics & Information Sciences An International Journal

http://dx.doi.org/10.18576/amis/190604

Advanced Algebraic Approaches to Soliton Solutions in Shallow Water Wave Equations

Mohamed A. Hafez ^{1,2}, Romana Ashraf³, Ali Akgül^{4,5*}, M. Qasymeh⁶, Shabbir Hussain⁷ and Farah Ashraf⁷

- ¹ Faculty of Engineering and Quantity Surviving, INTI International University Colleges, Nilai, Malaysia
- ² Faculty of Management, Shinawatra University, Pathum Thani, Thailand
- ³ Department of Computer Science & IT, The University of Lahore, Pakistan
- ⁴ Department of Electronics and Communication Engineering, Saveetha School of Engineering, SIMATS, Chennai, Tamilnadu, India
- ⁵ Department of Mathematics, Art and Science Faculty, Siirt University, 56100 Siirt, Turkey
- ⁶ Electrical and Computer Engineering Department, Abu Dhabi University, Abu Dhabi, United Arab Emirates
- ⁷ Department of Mathematics and Statistics, The University of Lahore, Lahore, Pakistan

Received: 15 Feb. 2025, Revised: 2 May 2025, Accepted: 19 Aug. 2025

Published online: 1 Nov. 2025

Abstract: In this paper, soliton solutions of the (1+1)-dimensional Boussinesq equation are studied with an improved scheme of the direct algebraic method. The Boussinesq equation as concerns the propagation of long waves in shallow water in addition to other physical systems, exhibit both dispersal and nonlinearity. With this new approach we obtain analytical soliton solutions and make a study of their characteristics. The obtained solutions are said in terms of positive forms and this is an improvement of the perception of the dynamics of the equation. The performance of the method is illustrated by calculation, and further use of the method in the related nonlinear systems is also discussed. This work adds to an ongoing research on the analysis of the solution of nonlinear partial differential equations and provides understanding on the solitons in the dispersive media. The obtained solutions as well have been depicted graphically for better understanding.

Keywords: Boussinesq equation, traveling wave solutions, new extended direct algebraic method for Boussinesq equation.

1 Introduction

Nonlinear evolution equations (NLEEs) as a class of equations are derived from the nonlinear sciences and those have important roles in analyze of the non-linear phenomenon. Another well preserved by self- reinforcement wave packet is referred to as soliton or solitary wave that moves at a constant velocity. The nonlinearity and dispersion of the medium are elimi- nated which in a result cause the formation of solitons. The class of weakly nonlinear dispersive PDEs which models physical systems has socalled solitons In non-linear PDEs, more striking phenomena exist than soli- ton solutions. A specific kind of a localized traveling wave solution of a nonlinear PDE which is immensely stable is known as a soliton [1–20]. Non-linear dispersive media allow wave pulses to propagate because of their dispersive effects as well as the dynamical balance between non-linearity [66].

It is used for analysis of long waves with small amplitude in shallow water and other physical processes in nonlinear lattice theory and plasma physics. In its (1+1)-dimensional form, the equation is expressed as:

$$Q_{tt} - Q_{xx} = \lambda Q_{xxxx} + (Q^2)_{xx}, \tag{1}$$

where Q(x,t) the wave profile and λ is a parameter that determines the scale of dispersion. The coordinates nonlinearity and dispersion in order to form soliton solutions i.e., localized increments that retain their form and propagate at a uniform velocity. The investigation of solitons in the framework of the Boussinesq equation is importance for the analysis of wave processes in different physical systems. Computational procedures of classical analysis to solve a system of nonlinear PDEs encounter difficulties in finding exact expressions. Thus, it is criti-

^{*} Corresponding author e-mail: aliakgul00727@gmail.com



cal to develop new analytical approaches that will help to study such systems.

The Boussinesq equation captures the interplay between dispersion and nonlinearity in wave propagation. The term λQ_{xxxx} â represents the dispersive effect, which tends to spread the wave out, while the term $(Q^2)_{xx}$ accounts for the nonlinear interaction, which can lead to wave steepening and the formation of solitons.

From a physical perspective of one and a half-dimensional Boussinesq equation, especially the newly derived solitons, one can easily see the complexities of the interaction between the dispersion and nonlinearity in wave motion. Using the new extended direct algebraic method we have been able to found types of traveling wave solutions which include singular soliton solutions, periodic soliton solutions, rational soliton solutions and both dark and bright soliton solutions. These stand for different wave phenomena which are possible in dispersive medium, for example, shallow water waves or nonlinear lattices. For example, the bright solitons correspond to the localized wave of elevation while the dark solitons describe localized drops of the wave profile. This should give a guarantee to the efficiency of the proposed method and the possibility to analyze the waves behaviors in different physical systems by the help of such opportunities to find soliton solutions with the use of different forms of initial functions. The results obtained in this work have an important implications for the analvsis of wave processes in the dispersive media and also in the fields such as mathematical biology, physics, chemistry and fluid dynamics where similar nonlinear partial differential equations are used for modeling of the complex phenomena. The accurate modeling of shallow water waves as addressed in this paper can directly support effective "water management and pollution remediation" and "freshwater and water quality" strategies.

A vast amount of work has been carried out in the last few decades studying the properties of nonlinear partial differential equations for the reason that they provide very ideal models of phenomena in various science and engineering disciplines. In fact, the exact solutions of the above equations help hand and glove in analyzing the various global physical phenomena and dynamism of the process. These kinds of solutions are required for interpreting the qualitative properties of a number of occurrences in various branches of the natural science. In the last few years much attention has been paid to attempt to find exact solutions by using nonlinear partial differential equation models. Since most nonlinear PDEs do not have closedform solutions, the study of such solutions whether they are travelling wave solutions or other classes of solutions are desirable for developing the understanding of these systems.

The methods for obtaining exact explicit solutions of nonlinear partial differential equations are the tanh-function

method [41], the exp-function method [42], the F-expansion method [43], Hirota's direct method [44], Kudryashov method [45], the modified extended direct algebraic method [64], the extended auxiliary equation method [47], modified method of simplest equation [48] and the new extended direct algebraic method [62]. Examples of the methods for solving nonlinear partial differential equations numerically are the finite element method [31, 32], finite volume method [33, 34], generalized finite difference method [35, 36], collocation method [37, 38] and Galerkin finite element method [39, 40].

2 Algorithm for New Extended Direct Algebraic Method

In this section, the algorithm for the new extended direct algebraic method is introduced, see [49,50]. In the following we will outline the main steps of our method.

Consider a general nonlinear PDE in the form

$$Q(u, u_x, u_y, u_z, u_w, u_t, u_{tt},) = 0, (2)$$

where Q is a polynomial function of its argument, and the subscripts denote partial derivatives. We seek its traveling wave solutions by using transformation wave

$$u(x, y, z, w, t) = U(\zeta), \qquad \zeta = ax + by + cz + dw + vt. \tag{3}$$

Substituting Eq. 3 into Eq. 2 yields a nonlinear ordinary differential equation

$$Q(u, u', u'', u''', \dots) = 0, (4)$$

where the prime denotes differentiation with respect to ζ . Let us consider that Eq.4 has a formal solution of the form

$$U(\zeta) = \sum_{j=0}^{n} H_j P^j(\zeta), \ H_n \neq 0, \tag{5}$$

where the H_j $(0 \le j \le n)$ are constants coefficients to be determined later, and n is a positive integer which is found by homogenous balancing principle between the highest nonlinear term and the highest derivative in Eq. 4 and $P(\zeta)$ satisfies the NODE Eq. 4 in the form of

$$P^{'}(\zeta) = \ln(A) \left(\Theta + \Psi P(\zeta) + \Gamma P^{2}(\zeta)\right), \quad A \neq (0,1), \quad (6)$$

where Θ , Ψ and Γ are constants. Some special solutions of the NODE are given in [49,50].



3 Application of the new extended direct algebraic method

In numerical models, water waves, and coastal engineering, the Boussinesq equation is a long-wavelength and weakly nonlinear approximation that is used to simulate water waves in shallow seas and harbours.

A Scottish engineer named John Scott Russell closely observed solitary waves. John Scott Russell's observation served as the basis for Joseph Boussinesq's approximation. Boussinesq's simulation of one-dimensional water waves in 1872 established that, in addition to the water depth, the vertical velocity is linear and the horizontal velocity is constant [27].

Considering (1+1) dimensional Boussinesq equation

$$Q_{tt} - Q_{xx} = \lambda Q_{xxxx} + \left(Q^2\right)_{xx}. (7)$$

where Q = Q(x,t) represents the wave envelope containing x as a spatial variable and t as a temporal variable. Here λ is an arbitrary constant.

When water waves of various wavelengths are related, this is referred to as a frequency dispersion phenomena; in the case of an infinitesimal wave amplitude, it is also referred to as a linear frequency dispersion. This makes the approximation accurate. Although waves can propagate in multiple directions according to the Boussinesq equation, it is more advantageous to take these into account. The Boussinesq equation utilizing a new extended direct algebraic technique for generating strong and reliable soltons. Now using the traveling wave transformation for this purpose

$$Q(x,t) = q(\zeta), \tag{8}$$

where $\zeta = kx + vt$,

$$Q(x,t) = q(kx + vt), (9)$$

substituting Eq. 9 into Eq. 7. We have the following NODE

$$v^{2}q''(\zeta) - k^{2}q''(\zeta) - k^{2}(q^{2}(\zeta))'' - \lambda k^{4}q^{i\nu}(\zeta) = 0, (10)$$

Integrating Eq. 10 twice and neglecting the constants of integration

$$(v^2 - k^2) q(\zeta) - k^2 q^2(\zeta) - \lambda k^4 q''(\zeta), \tag{11}$$

Balancing q'' with q^2 in Eq. 11 gives N=2. Thus, Eq. 11 has the formal solution

$$q(\zeta) = H_0 + H_1 P(\zeta) + H_2 P^2(\zeta), \tag{12}$$

substituting Eq. 12 along with Eq. 6 into Eq. 11 and setting the coefficients of all powers of P^i , $i=0,1\cdots$, to zero, we yield the a system of algebraic equations and solved this system with the aid of Maple or Mathematica, we obtained following values for constants;

Case 1

$$\begin{cases} H_0 = -6\lambda k^2 \Gamma \Theta \ln^2(A), \\ H_1 = -6\lambda k^2 \Psi \Gamma \ln^2(A), \\ H_2 = -6\lambda k^2 \Gamma^2 \ln^2(A), \\ v = \sqrt{1 + \lambda k^2 \Psi^2 - 4\lambda k^2 \Gamma \Theta} \ln(A). \end{cases}$$
(13)

Case 2

$$\begin{cases} H_0 = -\lambda k^2 \left(\Psi^2 + 2\Gamma \Theta \right) \ln^2(A), \\ H_1 = -6\lambda k^2 \Psi \Gamma \ln^2(A), \\ H_2 = -6\lambda k^2 \Gamma^2 \ln^2(A), \\ v = k\sqrt{1 - \lambda k^2 \Psi^2 + 4\lambda k^2 \Gamma \Theta} \ln(A). \end{cases}$$
(14)

We established following families of exact solutions using Eq.13 along with Eq.12 into Eq. 7 **Family1**. When $\Psi^2 - 4\Theta\Gamma < 0$ and $\Gamma \neq 0$, then the traveling wave solutions are given by



$$Q_{1,1}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \ln^2(A)\Psi\Gamma\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma}\tan_A\left(\frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}\zeta}{2}\right)\right] - 6\lambda k^2 \ln^2(A)\Gamma^2\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma}\tan_A\left(\frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}\zeta}{2}\right)\right]^2$$
(15)

$$Q_{2,1}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \Psi\Gamma \ln^2(A) \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \cot_A \left(\frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2}\zeta\right) \right] \\ - 6\lambda k^2 \ln^2(A)\Gamma^2 \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \cot_A \left(\frac{\sqrt{-(\Psi^2 - 4\Theta\Gamma)}}{2}\zeta\right) \right]^2$$
(16)

$$Q_{3,1}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A) \Gamma \Theta\right) - 6\lambda k^{2} \ln^{2}(A) \Psi \Gamma$$

$$\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^{2} - 4\Theta \Gamma)}}{2\Gamma} \left(\tan_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \pm \sqrt{pq} \sec_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \right) \right] - 6\lambda k^{2} \ln^{2}(A) \Gamma^{2}$$

$$\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^{2} - 4\Theta \Gamma)}}{2\Gamma} \left(\tan_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \pm \sqrt{pq} \sec_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \right) \right]^{2}$$
(17)

$$Q_{4,1}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A) \Gamma \Theta\right) - 6\lambda k^{2} \ln^{2}(A) \Psi \Gamma$$

$$\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^{2} - 4\Theta \Gamma)}}{2\Gamma} \left(-\cot_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \right) \right] - 6\lambda k^{2} \ln^{2}(A) \Gamma^{2}$$

$$\left[-\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^{2} - 4\Theta \Gamma)}}{2\Gamma} \left(-\cot_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(\sqrt{-(\Psi^{2} - 4\Theta \Gamma)} \zeta \right) \right) \right] \right]^{2}$$
(18)

$$Q_{5,1}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma$$

$$\left\{ -\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^2 - 4\Theta \Gamma)}}{4\Gamma} \left(\tan_A \left(\frac{1}{4} \sqrt{-(\Psi^2 - 4\Theta \Gamma)} \zeta \right) - \cot_A \left(\frac{1}{4} \sqrt{-(\Psi^2 - 4\Theta \Gamma)} \zeta \right) \right) \right\}$$

$$- 6\lambda k^2 \ln^2(A) \Gamma^2 \left\{ -\frac{\Psi}{2\Gamma} + \frac{\sqrt{-(\Psi^2 - 4\Theta \Gamma)}}{4\Gamma} \left(\tan_A \left(\frac{1}{4} \sqrt{-(\Psi^2 - 4\Theta \Gamma)} \zeta \right) - \cot_A \left(\frac{1}{4} \sqrt{-(\Psi^2 - 4\Theta \Gamma)} \zeta \right) \right) \right\}^2. \quad (19)$$

Family2. When $\Psi^2 - 4\Theta\Gamma > 0$ and $\Gamma \neq 0$, then the traveling wave solutions are given by

$$Q_{6,2}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \Psi\Gamma \ln^2(A) \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \tanh_A \left(\frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}\zeta}{2} \right) \right] \\ - 6\lambda k^2 \ln^2(A)\Gamma^2 \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \tanh_A \left(\frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}\zeta}{2} \right) \right]^2 \quad (20)$$

$$Q_{7,2}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \Psi\Gamma \ln^2(A) \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \coth_A\left(\frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}\zeta}{2}\right) \right] - 6\lambda k^2 \ln^2(A)\Gamma^2 \left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}}{2\Gamma} \coth_A\left(\frac{\sqrt{(\Psi^2 - 4\Theta\Gamma)}\zeta}{2}\right) \right]^2$$
(21)



$$\begin{split} Q_{8,2}\left(\zeta\right) &= \left(-6\lambda\,k^2\ln^2(A)\Gamma\,\Theta\right) - 6\lambda\,k^2\,\Psi\Gamma\,\ln^2(A) \\ &\left\{ -\frac{\Psi}{2\,\Gamma} + \frac{\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}}{2\Gamma}\left(-\tanh_A\left(\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}\,\zeta\right) \pm i\,\sqrt{p\,q}\,sech_A\left(\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}\,\zeta\right)\right)\right\} \\ &- 6\lambda\,k^2\ln^2(A)\,\Gamma^2\left\{ -\frac{\Psi}{2\,\Gamma} + \frac{\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}}{2\Gamma}\left(-\tanh_A\left(\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}\,\zeta\right) \pm i\,\sqrt{p\,q}\,sech_A\left(\sqrt{(\Psi^2 - 4\,\Theta\,\Gamma)}\,\zeta\right)\right)\right\}^2 \end{split} \tag{22}$$

$$Q_{9,2}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A)\Gamma\Theta\right) - 6\lambda k^{2} \ln^{2}(A)\Psi\Gamma$$

$$\left\{-\frac{\Psi}{2\Gamma} + \frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}}{2\Gamma}\left(-\coth_{A}\left(\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta\right) \pm \sqrt{pq} \operatorname{csch}_{A}\left(\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta\right)\right)\right\}$$

$$-6\lambda k^{2} \ln^{2}(A)\Gamma^{2}\left\{-\frac{\Psi}{2\Gamma} + \frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}}{2\Gamma}\left(-\coth_{A}\left(\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta\right) \pm \sqrt{pq} \operatorname{csch}_{A}\left(\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta\right)\right)\right\}^{2}$$
(23)

$$Q_{10,2}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A)\Gamma\Theta\right) - 6\lambda k^{2}\Psi\Gamma \ln^{2}(A)$$

$$\left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}}{4\Gamma}\left\{\tanh_{A}\left(\frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta}{4}\right) + \coth_{A}\left(\frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta}{4}\right)\right\}\right] - 6\lambda k^{2} \ln^{2}(A)\Gamma^{2}$$

$$\left[-\frac{\Psi}{2\Gamma} - \frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}}{4\Gamma}\left\{\tanh_{A}\left(\frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta}{4}\right) + \coth_{A}\left(\frac{\sqrt{(\Psi^{2} - 4\Theta\Gamma)}\zeta}{4}\right)\right\}\right]^{2}$$
(24)

Family3. When $\Theta \Gamma > 0$ and $\Psi = 0$, then the traveling wave solutions are given by

$$Q_{11,3}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \Psi \Gamma \ln^2(A) \left(\sqrt{\frac{\Theta}{\Gamma}} \tan_A \left(\sqrt{\Theta \Gamma} \zeta\right)\right) - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[\sqrt{\frac{\Theta}{\Gamma}} \tan_A \left(\sqrt{\Theta \Gamma} \zeta\right)\right]^2$$
(25)

$$Q_{12,3}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \Psi \Gamma \ln^2(A) \left(-\sqrt{\frac{\Theta}{\Gamma}} \cot_A \left(\sqrt{\Theta \Gamma} \zeta\right)\right) - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\sqrt{\frac{\Theta}{\Gamma}} \cot_A \left(\sqrt{\Theta \Gamma} \zeta\right)\right]^2$$
(26)

$$Q_{13,3}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A) \Gamma \Theta\right) - 6\lambda k^{2} \ln^{2}(A) \Psi \Gamma \left[\sqrt{\frac{\Theta}{\Gamma}} \left(\tan_{A} \left(2\sqrt{\Theta \Gamma}\zeta\right) \pm \sqrt{\frac{pq\Theta}{\Gamma}}\right) \sec_{A} \left(2\sqrt{\Theta \Gamma}\zeta\right)\right] - 6\lambda k^{2} \ln^{2}(A) \Gamma^{2} \left[\sqrt{\frac{\Theta}{\Gamma}} \left(\tan_{A} \left(2\sqrt{\Theta \Gamma}\zeta\right) \pm \sqrt{\frac{pq\Theta}{\Gamma}}\right) \sec_{A} \left(2\sqrt{\Theta \Gamma}\zeta\right)\right]^{2} \right]$$
(27)

$$Q_{14,3}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A) \Gamma \Theta\right) - 6\lambda k^{2} \ln^{2}(A) \Psi \Gamma \left[\sqrt{\frac{\Theta}{\Gamma}} \left(\cot_{A}\left(2\sqrt{\Theta \Gamma}\zeta\right) \pm \sqrt{\frac{pq\Theta}{\Gamma}}\right) \csc_{A}\left(2\sqrt{\Theta \Gamma}\zeta\right)\right] - 6\lambda k^{2} \ln^{2}(A) \Gamma^{2} \left[\sqrt{\frac{\Theta}{\Gamma}} \left(\cot_{A}\left(2\sqrt{\Theta \Gamma}\zeta\right) \pm \sqrt{\frac{pq\Theta}{\Gamma}}\right) \csc_{A}\left(2\sqrt{\Theta \Gamma}\zeta\right)\right]^{2} \right]$$
(28)

$$Q_{15,3}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \ln^2(A)\Psi\Gamma\left[1/2\sqrt{\frac{\Theta}{\Gamma}}\left(\tan_A\left(\frac{1}{2}\sqrt{\Theta\Gamma}\zeta\right) - \cot_A\left(\frac{1}{2}\sqrt{\Theta\Gamma}\zeta\right)\right)\right] - 6\lambda k^2 \ln^2(A)\Gamma^2\left[\frac{1}{2}\sqrt{\frac{\Theta}{\Gamma}}\left(\tan_A\left(\frac{1}{2}\sqrt{\Theta\Gamma}\zeta\right) - \cot_A\left(\frac{1}{2}\sqrt{\Theta\Gamma}\zeta\right)\right)\right]^2$$
(29)



Family4. When $\Theta \Gamma < 0$ and $\Psi = 0$, then the traveling wave solutions are given by

$$Q_{16,4}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left[-\sqrt{-\frac{\Theta}{\Gamma}} \tanh_A \left(\sqrt{-\Theta \Gamma} \zeta\right)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\sqrt{-\frac{\Theta}{\Gamma}} \tanh_A \left(\sqrt{-\Theta \Gamma} \zeta\right)\right]^2$$
(30)

$$Q_{17,4}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left[-\sqrt{-\frac{\Theta}{\Gamma}} \coth_A\left(\sqrt{-\Theta \Gamma}\zeta\right)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\sqrt{-\frac{\Theta}{\Gamma}} \coth_A\left(\sqrt{-\Theta \Gamma}\zeta\right)\right]^2$$
(31)

$$\begin{split} Q_{18,4}\left(\zeta\right) &= \left(-6\,\lambda\,k^2\ln^2(A)\varGamma\,\Theta\right) - 6\,\lambda\,k^2\ln^2(A)\,\Psi\varGamma\left[\sqrt{-\frac{\Theta}{\varGamma}}\left(-\tanh\left(2\sqrt{-\Theta\,\varGamma}\,\zeta\right)\right) \pm i\sqrt{pq}\,sech_A(2\sqrt{-\Theta\,\varGamma}\,\zeta)\right] \\ &\quad - 6\,\lambda\,k^2\ln^2(A)\,\varGamma^2\left[\sqrt{-\frac{\Theta}{\varGamma}}\left(-\tanh\left(2\sqrt{-\Theta\,\varGamma}\,\zeta\right)\right) \pm i\sqrt{pq}\,sech_A(2\sqrt{-\Theta\,\varGamma}\,\zeta)\right]^2 \end{split} \tag{32}$$

$$Q_{19,4}(\zeta) = \left(-6\lambda k^{2} \ln^{2}(A) \Gamma \Theta\right) - 6\lambda k^{2} \ln^{2}(A) \Psi \Gamma \left[\sqrt{-\frac{\Theta}{\Gamma}} \left(-\coth_{A}\left(2\sqrt{-\Theta\Gamma}\zeta\right) \pm \sqrt{pq} csch_{A}\left(2\sqrt{-\Theta\Gamma}\zeta\right)\right)\right] - 6\lambda k^{2} \ln^{2}(A) \Gamma^{2} \left[\sqrt{-\frac{\Theta}{\Gamma}} \left(-\coth_{A}\left(2\sqrt{-\Theta\Gamma}\zeta\right) \pm \sqrt{pq} csch_{A}\left(2\sqrt{-\Theta\Gamma}\zeta\right)\right)\right]^{2} \right]$$
(33)

$$Q_{20,4}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left[-\frac{1}{2} \sqrt{-\frac{\Theta}{\Gamma}} \left(\tanh_A \left(\frac{1}{2} \sqrt{-\Theta \Gamma} \zeta \right) + \coth_A \sqrt{-\Theta \Gamma} \zeta \right) \right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\frac{1}{2} \sqrt{-\frac{\Theta}{\Gamma}} \left(\tanh_A \left(\frac{1}{2} \sqrt{-\Theta \Gamma} \zeta \right) + \coth_A \sqrt{-\Theta \Gamma} \zeta \right) \right]^2 \right]$$
(34)

Family5: when $\Psi = 0$ and $\Gamma = \Theta$, then the singular periodic solutions are given by

$$Q_{21,5}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left[\tan_A(\Theta \zeta)\right]$$

$$-6\lambda k^2 \ln^2(A) \Gamma^2 \left[\tan_A(\Theta \zeta)\right]^2$$
 (35)

$$Q_{22,5}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma\left(\ln^2 A\right) \left[-\cot_A(\Theta \zeta)\right]$$

$$-6\lambda k^2 \ln^2(A) \Gamma^2\left(\ln^2 A\right) \left[-\cot_A(\Theta \zeta)\right]^2$$
 (36)

$$Q_{23,5}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left[\left(\tan_A \left(2\Theta\zeta\right) \pm \sqrt{pq}\right) \sec_A \left(2\Theta\zeta\right) \right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[\left(\tan_A \left(2\Theta\zeta\right) \pm \sqrt{pq}\right) \sec_A \left(2\Theta\zeta\right) \right]^2$$
(37)

$$Q_{24,5}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma\left(\ln^2 A\right) \left[-\left(\cot_A \left(2\Theta \zeta\right) \pm \sqrt{pq}\right) \csc_A \left(2\Theta \zeta\right)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\left(\cot_A \left(2\Theta \zeta\right) \pm \sqrt{pq}\right) \csc_A \left(2\Theta \zeta\right)\right]$$
(38)



$$Q_{25,5}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma\left(\ln^2 A\right) \left[\frac{1}{2} \left(\tan\left(\frac{1}{2}\Theta\zeta\right) - \cot\left(\frac{1}{2}\Theta\zeta\right)\right)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[\frac{1}{2} \left(\tan\left(\frac{1}{2}\Theta\zeta\right) - \cot\left(\frac{1}{2}\Theta\zeta\right)\right)\right]^2$$
(39)

Family6. When $\Psi = 0$ and $\Gamma = -\Theta$, then the traveling wave solutions are given by

$$Q_{26,6}\left(\zeta\right) = \left(-6\lambda\,k^2\ln^2(A)\Gamma\,\Theta\right) - 6\lambda\,k^2\ln^2(A)\Psi\Gamma\left(\ln^2A\right)\left[-\tanh_A\left(\Theta\zeta\right)\right] \\ - 6\lambda\,k^2\ln^2(A)\Gamma^2\left[-\tanh_A\left(\Theta\zeta\right)\right]^2 \quad (40)$$

$$Q_{27,6}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left[-\coth_A(\Theta \zeta)\right]$$

$$-6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\coth_A(\Theta \zeta)\right]^2$$
 (41)

$$Q_{28,6}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma\Theta\right) - 6\lambda k^2 \ln^2(A) \Psi\Gamma\left(\ln^2 A\right) \left[-\tanh_A\left(2\Theta\zeta\right) \pm i\sqrt{pq} sech_A\left(2\Theta\zeta\right)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\tanh_A\left(2\Theta\zeta\right) \pm i\sqrt{pq} sech_A\left(2\Theta\zeta\right)\right]^2$$
(42)

$$Q_{29,6}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma\left(\ln^2 A\right) \left[-\coth_A(2\Theta\zeta) \pm \sqrt{pq} \operatorname{csch}_A(2\Theta\zeta)\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\coth_A(2\Theta\zeta) \pm \sqrt{pq} \operatorname{csch}_A(2\Theta\zeta)\right]^2$$
(43)

$$\begin{aligned} \mathcal{Q}_{30,6}\left(\zeta\right) &= \left(-6\lambda\,k^2\ln^2(A)\Gamma\,\Theta\right) - 6\lambda\,k^2\ln^2(A)\,\Psi\Gamma\left(\ln^2A\right) \left[-\frac{1}{2}\left(\tanh_A\left(\frac{1}{2}\,\Theta\,\zeta\right) + \coth_A\left(\frac{1}{2}\,\Theta\,\zeta\right)\right) \right] \\ &- 6\lambda\,k^2\ln^2(A)\,\Gamma^2 \left[-\frac{1}{2}\left(\tanh_A\left(\frac{1}{2}\,\Theta\,\zeta\right) + \coth_A\left(\frac{1}{2}\,\Theta\,\zeta\right)\right) \right]^2 \end{aligned} \tag{44}$$

Family7. When $\Psi^2 = 4\Theta \Gamma$ and $\Gamma = -\Theta$, then the traveling wave solutions are given by

$$Q_{31,7}(\zeta) = \left(-6\lambda k^2 \ln^2(A)\Gamma\Theta\right) - 6\lambda k^2 \ln^2(A)\Psi\Gamma\left(\ln^2A\right) \left[-2\frac{\Theta\left(\Psi\zeta LnA + 2\right)}{\Psi^2\zeta LnA}\right] - 6\lambda k^2 \ln^2(A)\Gamma^2 \left[-2\frac{\Theta\left(\Psi\zeta LnA + 2\right)}{\Psi^2\zeta LnA}\right]^2$$
(45)

Family 8: When $\Psi = 0, \Theta = mk, (m \neq 0)$ and $\Gamma = k$, then

$$Q_{32,8}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left[A^{k\zeta} - m\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[A^{k\zeta} - m\right]^2$$
(46)

Family 9 . When $\Psi = \Gamma = 0$, then

$$Q_{33,9}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left[\Theta \zeta \ln(A)\right]$$

$$-6\lambda k^2 \ln^2(A) \Gamma^2 \left[\Theta \zeta \ln(A)\right]^2$$
 (47)

Family 10 . When $\Psi = \Theta = 0$ then

$$Q_{34,10}(\zeta) = \left(6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left[\frac{-1}{\Gamma \zeta \ln(A)}\right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[\frac{-1}{\Gamma \zeta \ln(A)}\right]^2$$
(48)

Family 11. When $\Psi \neq 0, \Theta = 0$, then the traveling wave solutions are given by

$$Q_{35,11}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left[-\frac{p\Psi}{\Gamma \left(\cosh_A(\Psi \zeta) - \sinh_A(\Psi \zeta) + p\right)} \right] - 6\lambda k^2 \ln^2(A) \Gamma^2 \left[-\frac{p\Psi}{\Gamma \left(\cosh_A(\Psi \zeta) - \sinh_A(\Psi \zeta) + p\right)} \right]^2$$
(49)

$$Q_{36,11}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6\lambda k^2 \ln^2(A) \Psi \Gamma \left(\ln^2 A\right) \left\{ -\frac{\Psi(\sinh_A(\Psi \zeta) + \cosh_A(\Psi \zeta))}{\Gamma\left(\sinh_A(\Psi \zeta) + \cosh_A(\Psi \zeta) + q\right)} \right\} \\ - 6\lambda k^2 \ln^2(A) \Gamma^2 \left\{ -\frac{\Psi(\sinh_A(\Psi \zeta) + \cosh_A(\Psi \zeta))}{\Gamma\left(\sinh_A(\Psi \zeta) + \cosh_A(\Psi \zeta) + q\right)} \right\}^2$$
(50)

Family 12: When $\Psi = l, \Gamma = m \, (m \neq 0)$, and $\Theta = 0$ then the rational solution is given by

$$Q_{37,12}(\zeta) = \left(-6\lambda k^2 \ln^2(A) \Gamma \Theta\right) - 6lk^2 \Psi \Gamma \ln^2(A) \left(\ln^2 A\right) \left\{ -\frac{pA^{l\zeta}}{q - mpA^{l\zeta}} \right\}$$

$$-6lk^2 \Gamma^2 \ln^2(A) \left\{ -\frac{pA^{l\zeta}}{q - mpA^{l\zeta}} \right\}^2. \quad (51)$$

where ζ is an independent variable, p and q are arbitrary constants greater than zero and called deformation parameters.

4 The graphical representation




Fig. 1: The 3-D and 2-D graphs of $q_1(\zeta)$ given by Eq. 15.






Fig. 2: The 3-D and 2-D graphs of $q_2(\zeta)$ given by Eq. 16.




Fig. 3: The 3-D and 2-D graphs of $q_3(\zeta)$ given by Eq. 17.




Fig. 4: The 3-D and 2-D graphs of $q_4(\zeta)$ given by Eq. 18.




Fig. 5: The 3-D and 2-D graphs of $q_5(\zeta)$ given by Eq. 19.




Fig. 6: The 3-D and 2-D graphs of $q_6(\zeta)$ given by Eq. 20.




Fig. 7: The 3-D and 2-D graphs of $q_7(\zeta)$ given by Eq. 21.




Fig. 8: The 3-D and 2-D graphs of $q_8(\zeta)$ given by Eq. 22.




Fig. 9: The 3-D and 2-D graphs of $q_9(\zeta)$ given by Eq. 23.




Fig. 10: The 3-D and 2-D graphs of $q_{10}(\zeta)$ given by Eq. 24.




Fig. 11: The 3-D and 2-D graphs of $q_{11}(\zeta)$ given by Eq. 25.




Fig. 12: The 3-D and 2-D graphs of $q_{12}(\zeta)$ given by Eq. 26.




Fig. 13: The 3-D and 2-D graphs of $q_{13}(\zeta)$ given by Eq. 27.






Fig. 14: The 3-D and 2-D graphs of $q_{14}(\zeta)$ given by Eq. 28.




Fig. 15: The 3-D and 2-D graphs of $q_{15}(\zeta)$ given by Eq. 29.




Fig. 16: The 3-D and 2-D graphs of $q_{16}(\zeta)$ given by Eq. 30.




Fig. 17: The 3-D and 2-D graphs of $q_{20}(\zeta)$ given by Eq. 34.




Fig. 18: The 3-D and 2-D graphs of $q_{21}(\zeta)$ given by Eq. 35.




Fig. 19: The 3-D and 2-D graphs of $q_{26}(\zeta)$ given by Eq. 40.






Fig. 20: The 3-D and 2-D graphs of $q_{27}(\zeta)$ given by Eq. 41.




Fig. 21: The 3-D and 2-D graphs of $q_{28}(\zeta)$ given by Eq. 42.




Fig. 22: The 3-D and 2-D graphs of $q_{29}(\zeta)$ given by Eq. 43.




Fig. 23: The 3-D and 2-D graphs of $q_{30}(\zeta)$ given by Eq. 44.




Fig. 24: The 3-D and 2-D graphs of $q_{31}(\zeta)$ given by Eq. 45.




Fig. 25: The 3-D and 2-D graphs of $q_{35}(\zeta)$ given by Eq. 41.




Fig. 26: The 3-D and 2-D graphs of $q_{37}(\zeta)$ given by Eq. 51.



5 Discussion and Conclusion

In this study, we have applied a new extended direct algebraic method to obtain soliton solutions of the (1+1)dimensional Boussinesq equation. The derived solutions are presented in explicit forms, showcasing the method's capability to handle complex nonlinear PDEs. Our results contribute to the understanding of soliton dynamics in dispersive media and provide a foundation for future research in related areas. The success of this method in finding exact solutions highlights its potential as a valuable tool in the study of nonlinear wave equations, with implications across various physical and engineering disciplines. By the application of new extended direct algebraic method we have obtained a series of traveling wave solutions namely: singular solitons, periodic solitons, rational solitons, dark and bright soliton solutions for (1+1)-Boussinesq equation. The proposed method is straight forward and more powerful in constructing exact traveling wave solutions of NLPDE's. It can also be applied to other nonlinear partial differential appearing in mathematical biology, physics, chemistry, fluid mechanics and many other fields.

References

- [1] STR. Rizvi, AR. Seadawy, F. Ashraf, M. Younis, H. Iqbal, and D. Baleanu, 2020. Lump and interaction solutions of a geophysical Korteweg de Vries equation. Results in Physics, 19: 103661.
- [2] Farrah Ashraf, Aly R.Seadawy, Syed T.R.Rizvi, Kashif Ali, and M. Aamir Ashraf, 2022. Multi-wave, M-shaped rational and interaction solutions for fractional nonlinear electrical transmission line equation. Journal of Geometry and Physics, 177: 104503.
- [3] Rizvi. STR, Seadawy. AR, Ashraf. F, Younis .M, Iqbal. H, and Baleanu. D, 2019. Kinky breathers, W-shaped and multi-peak solitons interaction in (2+1)-dimensional nonlinear Schrodinger equation with Kerr law of nonlinearity. Eur. Phys. J. Plus, 134: 1-10.
- [4] Romana Ashraf, Faiza Amanat, Farah Ashraf, Soliton solutions for the (4+1)-dimensional Fokas equation using integration techniques, Alexandria Engineering Journal, 107 (2024) 61-72.
- [5] Ali Akgül, Saliha Manzoor, Farrah Ashraf, Romana Ashraf, Exact solutions of the (2+1)-dimensional Zoomeron model arising in nonlinear optics via mapping method, Optical and Quantum Electronics, 56 (7) (2024) 1207.
- [6] Shabbir Hussain, Muhammad Sajid Iqbal, Mustafa Bayram, Romana Ashraf, Mustafa Inc, Shahram Rezapour, Muhammad Akhtar Tarar, Optical soliton solutions in a distinctive class of nonlinear Schrödinger's equation with cubic, quintic, septic, and nonic nonlinearities, Optical and Quantum Electronics, 56 (6) (2024) 1066.
- [7] Mustafa Inc, Shabbir Hussain, Ali Hasan Ali, Muhammad Sajid Iqbal, Romana Ashraf, Muhammad Akhtar Tarar, Muhammad Adnan, Analyzing solitary wave solutions of the nonlinear Murray equation for blood flow in vessels with non-uniform wall properties, Scientific Reports, 10588 (2024).

- [8] F. Ashraf, R. Ashraf, A. Akgül, Travelling waves solutions of Hirota-Ramani equation by modified extended direct algebraic method and new extended direct algebraic method, International Journal of Modern Physics B, (2024).
- [9] R. Ashraf, S. Hussain, M. S. Iqbal, The extended Fan's sub-equation method and its application to nonlinear Schrödinger equation with saturable nonlinearity, Results in Physics, (2023).
- [10] S. Hussain, M. S. Igbal, R. Ashraf, M. Inc, M. A. Tarar, Exploring nonlinear dispersive waves in a disordered medium: an analysis using ?6 model expansion method, Optical and Quantum Electronics, 55 (7) (2023) 651.
- [11] S. Hussain, M. S. Iqbal, R. Ashraf, M. Inc, M. A. Tarar, Quantum analysis of nonlinear optics in Kerr affected saturable nonlinear media and multiplicative noise: a path to new discoveries, Optical and Quantum Electronics, 55 (7) (2023) 578.
- [12] R. Ashraf, F. Ashraf, A Akgül, S Ashraf, B Alshahrani, M Mahmoud, Some new soliton solutions to the (3+1)dimensional generalized KdV-ZK equation via enhanced modified extended tanh-expansion approach, Alexandria Engineering Journal, 69 (2023) 303-309.
- [13] S.T. R. Rizvi, Aly R. Seadawy, Sarfaraz Ahmed, R. Ashraf, Lax pair, Darboux transformation, Weierstrass-Jacobi elliptic and generalized breathers along with soliton solutions for Benjamin Bona Mahony equation, International Journal of Modern Physics B, (2023) 2350233.
- [14] Aly R. Seadawy, S.T. R. Rizvi, Tahira Batool, R. Ashraf, Study of Sasa-Satsuma dynamical system for Kuznetsov-Ma and generalized breathers, lump, periodic and rogue wave solutions, International Journal of Modern Physics B, (2023) 2350181.
- [15] Farrah Ashraf, Tehsina Javeed, Romana Ashraf, Amina Rana, Ali Akgul, Some new soliton solution to the higher dimensional burger-huxley and shallow water waves equation with couple of integration architectonic, Results in Physics, 43 (2022) 106048.
- [16] S.T. R. Rizvi, Aly R. Seadawy, R. Ashraf, Propagation of chirped periodic and solitary waves for the coupled nonlinear Schrödinger equation in two core optical fibers with parabolic law with weak non-local nonlinearity, Optical and Quantum Electronics, 545 (2022).
- [17] S. Ahmed, R. Ashraf, Aly R. Seadawy, S.T. R. Rizvi, M. Younis, Ali Althobaiti, Ahmed M. El-Shehawi, Lump, multi-wave, Kinky breathers, interactional solutions, and stability analysis for general (2+1)-th dispersionless Dym equation, Results in Physics, 25 (2021) 104160.
- [18] R. Ashraf, M.O. Ahmad, M. Younis, K.U. Tariq, K. Ali, S.T.R. Rizvi: Dipole and combo solitons in DWDM systems, Optik, 158 (2018) 1073-1079.
- [19] R. Ashraf, M.O. Ahmad, M. Younis, K. Ali, S.T.R. Rizvi: Dipole and Gausson soliton for ultrashort laser pulse with high order dispersion, Superlattices and Microstructures, 109 (2017) 504-510.
- [20] S.T.R. Rizvi, S. Bashir, M. Younis, R. Ashraf, M.O. Ahmad: Ahmad: Exact soliton of (2+1)-dimensional fractional Schrödinger equation, Superlattices and Microstructures, 107 (2017) 234-239.
- [21] Alkhidhr, H. A., & Abdelrahman, M. A. (2021). Wave structures to the three coupled nonlinear Maccaris systems in plasma physics. Results in Physics, 105092.



- [22] Ziane, D., Hamdi Cherif, M., Baleanu, D., & Belghaba, K. (2020). Non-Differentiable Solution of Nonlinear Biological Population Model on Cantor Sets. Fractal and Fractional, 4(1), 5.
- [23] Younis, M., Sulaiman, T. A., Bilal, M., Rehman, S. U., & Younas, U. (2020). Modulation instability analysis, optical and other solutions to the modified nonlinear Schrodinger equation. Communications in Theoretical Physics, 72(6), 065001.
- [24] Younas, U., Seadawy, A. R., Younis, M., & Rizvi, S. T. R. (2020). Dispersive of propagation wave structures to the Dullin-Gottwald-Holm dynamical equation in a shallow water waves. Chinese Journal of Physics, 68, 348-364.
- [25] Kim, H., Bae, J. H., & Sakthivel, R. (2014). Exact travelling wave solutions of two important nonlinear partial differential equations. Zeitschrift fur Naturforschung A, 69(3-4), 155-162.
- [26] Paredes, A., Olivieri, D. N., & Michinel, H. (2020). From optics to dark matter: A review on nonlinear Schrodinger-Poisson systems. Physica D: Nonlinear Phenomena, 403, 132301.
- [27] Yomba, E. (2005). The extended Fan's sub-equation method and its application to KdV-MKdV, BKK and variant Boussinesq equations. Physics Letters A, 336(6), 463-476.
- [28] Batool, F., & Akram, G. (2017). Application of extended Fan sub-equation method to (1+ 1)-dimensional nonlinear dispersive modified Benjamin-Bona-Mahony equation with fractional evolution. Optical and Quantum Electronics, 49(11), 1-9.
- [29] El-Wakil, S. A., & Abdou, M. A. (2008). The extended Fan sub-equation method and its applications for a class of nonlinear evolution equations. Chaos, Solitons & Fractals, 36(2), 343-353.
- [30] Cheemaa, N., & Younis, M. (2016). New and more exact traveling wave solutions to integrable (2+ 1)-dimensional Maccari system. Nonlinear Dynamics, 83(3), 1395-1401.
- [31] Lakkis, O., & Pryer, T. (2013). A finite element method for nonlinear elliptic problems. SIAM Journal on Scientific Computing, 35(4), A2025-A2045.
- [32] Yan, S., & Jin, J. M. (2015). Theoretical formulation of a time-domain finite element method for nonlinear magnetic problems in three dimensions. Progress In Electromagnetics Research, 153, 33-55.
- [33] Carrillo, J. A., Chertock, A., & Huang, Y. (2015). A finite-volume method for nonlinear nonlocal equations with a gradient flow structure. Communications in Computational Physics, 17(1), 233-258.
- [34] Murakawa, H. (2017). A linear finite volume method for nonlinear cross-diffusion systems. Numerische Mathematik, 136(1), 1-26.
- [35] Gavete, L., Urea, F., Benito, J. J., Garca, A., Urea, M., & Salete, E. (2017). Solving second order non-linear elliptic partial differential equations using generalized finite difference method. Journal of Computational and Applied Mathematics, 318, 378-387.
- [36] Urea, F., Gavete, L., Garcia, A., Benito, J. J., & Vargas, A. M. (2019). Solving second order non-linear parabolic PDEs using generalized finite difference method (GFDM). Journal of Computational and Applied Mathematics, 354, 221-241.
- [37] Oru. (2020). An efficient wavelet collocation method for nonlinear two-space dimensional Fisher-Kolmogorov-Petrovsky-Piscounov equation and two-space dimensional

- extended Fisher-Kolmogorov equation. Engineering with Computers, 36(3), 839-856.
- [38] Wu, H., Wang, Y., & Zhang, W. (2018). Numerical solution of a class of nonlinear partial differential equations by using barycentric interpolation collocation method. Mathematical Problems in Engineering, 2018.
- [39] Ucar, Y., Karaagac, B., & Esen, A. (2015). A new approach on numerical solutions of the Improved Boussinesq type equation using quadratic B-spline Galerkin finite element method. Applied Mathematics and Computation, 270, 148-155.
- [40] Li, M., Huang, C., & Wang, P. (2017). Galerkin finite element method for nonlinear fractional Schrodinger equations. Numerical Algorithms, 74(2), 499-525.
- [41] Abdusalam, H. A. (2005). On an improved complex tanhfunction method. International Journal of Nonlinear Sciences and Numerical Simulation, 6(2), 99-106.
- [42] Aminikhah, H., Moosaei, H., & Hajipour, M. (2010). Exact solutions for nonlinear partial differential equations via Expfunction method. Numerical Methods for Partial Differential Equations, 26(6), 1427-1433.
- [43] Zhao, Y. M. (2013). F-expansion method and its application for finding new exact solutions to the Kudryashov-Sinelshchikov equation. Journal of Applied Mathematics, 2013.
- [44] Wazwaz, A. M. (2008). The Hirota s direct method for multiple-soliton solutions for three model equations of shallow water waves. Applied Mathematics and Computation, 201(1-2), 489-503.
- [45] Ryabov, P. N., Sinelshchikov, D. I., & Kochanov, M. B. (2011). Application of the Kudryashov method for finding exact solutions of the high order nonlinear evolution equations. Applied Mathematics and Computation, 218(7), 3965-3972.
- [46] Younis, M., Seadawy, A. R., Baber, M. Z., Yasin, M. W., Rizvi, S. T., & Iqbal, M. S. Abundant solitary wave structures of the higher dimensional Sakovich dynamical model. Mathematical Methods in the Applied Sciences.
- [47] Jiong, S. (2003). Auxiliary equation method for solving nonlinear partial differential equations. Physics Letters A, 309(5-6), 387-396.
- [48] Vitanov, N. K. (2011). On modified method of simplest equation for obtaining exact and approximate solutions of nonlinear PDEs: the role of the simplest equation. Communications in Nonlinear Science and Numerical Simulation, 16(11), 4215-4231.
- [49] Mirhosseini-Alizamini, S., Rezazadeh, H., Srinivasa, K. & Bekir, A. New closed form solutions of the new coupled Konnoâ Oono equation using the new extended direct algebraic method. *Pramana*. 94, 1-12 (2020)
- [50] Rezazadeh, H., Tariq, H., Eslami, M., Mirzazadeh, M. & Zhou, Q. New exact solutions of nonlinear conformable time-fractional Phi-4 equation. *Chinese Journal Of Physics*. 56, 2805-2816 (2018)
- [51] Khan, A., Seadawy, A. R., & Nadeem, M. (2021). Soliton solutions for the fractional complex Ginzburg Landau equation using the Kudryashov method. Applied Mathematics & Information Sciences, 15(1), 73-79.
- [52] Nisar, K. S., Kilicman, A., & Agarwal, P. (2018). Optical soliton solutions for the perturbed Gerdjikov Ivanov equation with Kerr law nonlinearity. Applied Mathematics & Information Sciences, 12(6), 1233-1240.



- [53] Baleanu, D., Jajarmi, A., & Asad, J. H. (2018). New aspects of the fractional optimal control problems: Mittag-Leffler stability and exact solutions. Progress in Fractional Differentiation and Applications, 4(3), 161-172.
- [54] Gómez-Aguilar, J. F., & Atangana, A. (2017). Fractional Schrödinger equation with non-local and non-singular kernel: A model for nonlinear waves in complex media. Progress in Fractional Differentiation and Applications, 3(2), 85-98.
- [55] El-Depsy, A., & Al-Marzoug, S. M. (2021). Optical Solitons in Nonlinear Thin-Film Waveguides: A Theoretical Approach. International Journal of Thin Film Science and Technology, 10(2), 45-53.
- [56] Rahman, M. M., & Hossain, M. A. (2020). Mathematical Modeling of Heat Transfer in Nanoscale Thin Films Using Fractional PDEs. International Journal of Thin Film Science and Technology, 9(3), 87-95.
- [57] Rahman, M., & Basak, S. (2022). Statistical Properties of Soliton Solutions in the (3+1)-Dimensional Nonlinear Schrödinger Equation. Journal of Statistics Applications & Probability, 11(1), 45-58.
- [58] Alzaatreh, A., & Al-Labadi, L. (2021). A Statistical Approach to Modeling Nonlinear Wave Phenomena with Applications to Soliton Solutions. Journal of Statistics Applications & Probability, 10(2), 215-225.
- [59] Iqbal, M. S. (2011). Solutions of boundary value problems for nonlinear partial differential equations by fixed point methods, T.U. Graz library.
- [60] Rehman, H., Seadawy, A., Younis, M., Rizvi, S., Anwar, I., Baber, M. & Althobaiti, A. Weakly nonlinear electron-acoustic waves in the fluid ions propagated via a (3+ 1)-dimensional generalized Kortewegâ de-Vriesâ Zakharovâ Kuznetsov equation in plasma physics. *Results In Physics*. 33 pp. 105069 (2022)
- [61] Yakada, S., Depelair, B., Betchewe, G. & Doka, S. Miscellaneous new traveling waves in metamaterials by means of the new extended direct algebraic method. *Optik.* 197 pp. 163108 (2019)
- [62] Mirhosseini-Alizamini, S., Rezazadeh, H., Eslami, M., Mirzazadeh, M. & Korkmaz, A. New extended direct algebraic method for the Tzitzica type evolution equations arising in nonlinear optics. *Computational Methods For Differential Equations*. 8, 28-53 (2020)
- [63] Vahidi, J., Zabihi, A., Rezazadeh, H. & Ansari, R. New extended direct algebraic method for the resonant nonlinear SchrA¶dinger equation with Kerr law nonlinearity. *Optik.* 227 pp. 165936 (2021)
- [64] Younis, M., Seadawy, A., Baber, M., Husain, S., Iqbal, M., Rizvi, S. & Baleanu, D. Analytical optical soliton solutions of the Schrodinger-Poisson dynamical system. *Results In Physics*. 27 pp. 104369 (2021)
- [65] Kaplan, M., Akbulut, A. Raza, N. Research on sensitivity analysis and traveling wave solutions of the (4+ 1)-dimensional nonlinear Fokas equation via three different techniques. *Physica Scripta*. 97, 015203 (2022)
- [66] Alzkari T., Wei Sin Koh, Optimal Solutions of the Time-Fractional Wave Models, *International Journal of Analysis* and Applications. 23, 129 (2025)



Mohamed Ahmed Hafez earned his Ph. D. degree in Civil Engineering from University of Malaya. an Assoc. Prof. He is at the Department of Civil Engineering, INTI International University. He has published over 55 papers in journals. His research

interests are focused on Dam risk, slope stability and soft ground improvement.



Romana Ashraf is an assistant professor University at The of Lahore, Department of Computer Science & IT. She is also working as Teaching Area InCharge (TAI) in the respective department. Her research areas are Nonlinear Partial Differential Equations,

Soliton Theory, Numerical Methods, Mathematical Modeling, and Fractional Differential Equations. She has published 21 research papers in internationally reputed journals, which have been extensively cited by researchers both nationally and internationally, with over 281 citations. She has a strong research profile, reflected by an h-index of 9 and an i10-index of 9. She has received the Research Productivity Award 2023 by The University of Lahore, Lahore.



Ali Akgül is a full professor in Siirt university, Faculty of Art and Science, Department of Mathematics. He is the head of the Mathematics department. His research areas are Fractional Differential Equations, Numerical Methods, Partial Differential Equations.

Mathematical Modeling and Functional Analysis. He made a big contribution on fractional calculus and numerical methods. He has more than 700 research papers in very good journals. He has given many talks as an invited speaker in many international conferences. He opened many special issues in very good journals. He is among the World's Top 2% Scientists by Stanford University in 2021, 2022, 2023 and 2024. He got OBADA prize in 2022 (Young Distinguished Researchers).





Montasir Qasymeh received the Ph.D. degree in electrical engineering from Dalhousie University, Halifax, Canada, in 2010, and completed a postdoctoral fellowship at the University of Ottawa, Canada, in 2011. He is currently the Associate Provost for Research,

Innovation, and Academic Development at Abu Dhabi University (ADU), United Arab Emirates, where he also serves as a Professor of Electrical Engineering. Dr. Qasymeh has played a transformative role in advancing ADU's research and innovation agenda. He led the establishment of the MENA region's first quantum computing laboratory, two interdisciplinary research institutes, and the Abu Dhabi Graphene Center. He has also developed robust frameworks to support industry-academia collaboration, startup incubation, and intellectual property commercialization. His research interests include quantum systems, nanoplasmonics, and terahertz wave engineering, with a particular focus on using two-dimensional graphene materials for the design of quantum and terahertz devices. He has authored and co-authored more than 70 peer-reviewed publications, including collaborative work on plasmonic sensors and devices. A prolific inventor, Dr. Qasymeh holds 17 U.S. patents and serves on the editorial board of a leading multidisciplinary journal. He has organized and chaired several high-impact conferences, including serving as General Co-Chair of the International Conference on Electrical, Communication, and Computer Engineering (ICEET) in 2023 and 2024, and as Subcommittee Chair for Quantum Science and Technology at PIERS 2025. Dr. Qasymeh is a frequent invited speaker at global scientific forums and actively contributes to international research and development initiatives.



Shabbir Hussain is a PhD scholar in the Department of Mathematics, University of Lahore, under the supervision of Dr. Romana Ashraf. His research include areas nonlinear partial differential equations, bifurcation theory, chaos analysis, soliton dynamics,

and analytical solution methods for complex nonlinear systems. He has more than 10 research papers in very good journals. He has given many talks as invited speaker in many national conferences.



Farrah Ashraf is an assistant professor at The University of Lahore, Lahore, Department of Mathematics and Statistics. Her research areas are Nonlinear Partial Differential Equations, Soliton Theory, Numerical Methods, Mathematical Modeling, Biomathematics

and Fractional Differential Equations. She has published 21 research papers in internationally reputed journals, which have been extensively cited by researchers both nationally and internationally, with over 326 citations. She has a strong research profile, reflected by an h-index of 8 and an i10-index of 7. She has received the Research Productivity Award 2023 by The University of Lahore, Lahore.