Applied Mathematics & Information Sciences An International Journal

http://dx.doi.org/10.18576/amis/200103

Analytical Exploration of the (4+1)-Dimensional Fokas Equation: Exact Traveling Wave Solutions

Mohamed A. Hafez ¹, Romana Ashraf², Ali Akgül^{3,4,*}, M. Qasymeh⁵, Shabbir Hussain⁶ and Farah Ashraf⁶

- ¹ Faculty of Engineering and Quantity Surviving, INTI International University Colleges, Nilai, Malaysia
- ² Department of Computer Science & IT, The University of Lahore, Pakistan
- ³ Department of Electronics and Communication Engineering, Saveetha School of Engineering, SIMATS, Chennai, Tamilnadu, India
- ⁴ Department of Mathematics, Art and Science Faculty, Siirt University, 56100 Siirt, Turkey
- ⁵ Electrical and Computer Engineering Department, Abu Dhabi University, Abu Dhabi, United Arab Emirates
- ⁶ Department of Mathematics and Statistics, The University of Lahore, Lahore, Pakistan

Received: 15 Mar. 2025, Revised: 2 Jun. 2025, Accepted: 19 Aug. 2025

Published online: 1 Jan. 2026

Abstract: This research paper aims at obtaining closed-form solutions for the traveling wave solutions of the (4+1)-dimensional Fokas equation. To solve this difficult system of nonlinear partial differential equations, we will use a new improved direct algebraic method. The given technique can be seen to work well for the purpose of converting the governing equation into a set of algebraic equations from where one can extract various traveling wave solutions. Here our results concern different forms of the solutions as solitons, periodic and rational functions. The obtained solutions are important for understanding the physical processes which are characterized by this equation and may be of use in Fluid mechanics, Nonlinear optics, Plasma physics and other areas.

Keywords: (4+1)-dimensional Fokas equation, traveling wave solutions, new extended direct algebraic method, bright soliton, dark soliton.

1 Introduction

Nonlinear evolution equations (NLEEs) as a class of equations are derived from the nonlinear sciences and those have important roles in analyze of the nonlinear preserved phenomenon. Another well self-reinforcement wave packet is referred to as soliton or solitary wave that moves at a constant velocity. The nonlinearity and dispersion of the medium are eliminated which in a result cause the formation of solitons. The class of weakly nonlinear dispersive PDEs which models physical systems has so-called solitons In nonlinear PDEs, more striking phenomena exist than soliton solutions. A specific kind of a localized traveling wave solution of a nonlinear PDE which is immensely stable is known as a soliton [1-20]. Nonlinear Partial differential equations (NPDEs) are a powerful and indispensable tool of modern mathematical modeling, applied for the description of many sophisticated phenomena in science and engineering. Evaluating these equations is at the heart of reproducing highly complex behaviors and remains

instrumental in developing theory and technology from physics and fluid mechanics to biological and financial systems. One of the issues arising with the study of NPDEs is the question of determining exact solutions to these equations, which is essential for determining the physical processes going on and for modeling many systems described by such equations. Traditionally, getting analytical solutions to NPDEs has been quite hard and even to date, some problems are vet to be solved. Since these equations are mutually dependent and quite elaborate at most times, simple solutions are rarely possible, which calls for enhanced methods. Much has been done in this area in the recent past perhaps due to the increase in computational power and availability of better algorithms. These have made the solution of NPDEs possible and have created opportunities for applying solutions previously pipe dreams. The sophisticated symbolic computation systems like Maple, Mathematica, MATLAB, etc. have gone further in encouraging the researchers by providing more efficient instruments to analyze broader areas of solution techniques. These tools

^{*} Corresponding author e-mail: aliakgul00727@gmail.com



have enabled one to obtain results that hitherto had been out of the reach of the researcher working on NPDEs. It enriches the existing research on exact solutions of NPDEs by presenting a new method suitable for a certain type of such equations. Thus, using the potential of the current methods of computational mathematics, this work intends to contribute to the development of the existing theoretical material and provide a new look at the behavior of the given complex systems that are characterized by NPDEs. The conclusions drawn in this paper contribute not only to the development of knowledge on these systems but also to the further research done in the fascinating and most diverse field.

The techniques for finding exact analytical solutions of nonlinear partial differential equations are the tanh function technique [41], the exp function technique [42], the F-expansion method [43], Hirota-s direct method [44], Kudryashov technique [45], the modified extended direct algebraic method [64], the extended auxiliary equation technique [47], the modified technique of the simplest equation [48] and the new extended direct algebraic technique [62]. Some of the methods adapted for numerical solution of the nonlinear partial differential equations are finite element method [31, 32], finite volume method [33, 34], generalized finite difference method [35, 36], collocation method [37, 38], Galerkin finite element method [39, 40].

In this paper, we are interested in applying the new extended direct algebraic method for solving the presented model trying to reveal their dynamics. The organization of the paper is as follows: in section 2, the algorithm for applying the new extended direct algebraic method is introduced with its main steps revealing different types of solutions. In section 3, the application of the presented method on the (4+1)-dimensional Fokas equation is illustrated along with the graphical representations of the obtained solutions. Finally, section 6 present the discussion of the obtained results along with a conclusion of the present and possible future work.

2 Algorithm for New Extended Direct Algebraic Method

In this section, the algorithm for the new extended direct algebraic method is introduced, see [49, 50]. In the following we will outline the main steps of our method. Consider a general nonlinear PDE in the form

$$Q(u, u_x, u_y, u_z, u_w, u_t, u_{tt}, \dots) = 0,$$
(1)

where Q is a polynomial function of its argument, and the subscripts denote partial derivatives. We seek its traveling wave solutions by using transformation wave

Q(u, u', u'', u''', ...) = 0,

Substituting Eq. 2 into Eq. 1 yields a nonlinear ordinary

differential equation

$$Q(u, u', u'', u''', \dots) = 0, (3)$$

where the prime denotes differentiation with respect to ζ . Let us consider that Eq.3 has a formal solution of the form

$$U(\zeta) = \sum_{j=0}^{n} \eta_j P^j(\zeta), \quad \eta_n \neq 0, \tag{4}$$

where the η_i ($0 \le j \le n$) are constants coefficients to be determined later, and n is a positive integer which is found by homogenous balancing principle between the highest nonlinear term and the highest derivative in Eq. 3 and $P(\zeta)$ satisfies the NODE Eq. 3 in the form of

$$P'(\zeta) = \ln A \left(\Psi + \Theta P(\zeta) + \Omega P^2(\zeta) \right), \quad A \neq (0, 1), \quad (5)$$

where Ψ , Θ and Ω are constants. Some special solutions of the NODE are given in [49, 50].

3 Application of the new extended direct algebraic method

Considering (4+1) – dimensional Fokas equation [65]

$$4u_{tx} - u_{xxxy} + u_{xyyy} + 12u_xu_y + 12uu_{xy} - 6u_{zw} = 0.$$
 (6)

u: dependent variable representing a physical quantity (e.g., wave amplitude, fluid velocity) x, y, z, t: independent variables representing spatial coordinates and time, respectively Making a wave transformation

$$u(x, y, z, w, t) = U(\zeta), \tag{7}$$

with $\zeta = ax + by + cz + dw + vt$.

Plugging Eq.7 into 6, then resulting nonlinear ordinary differential equation is:

$$(4av - 6cd)U''(\zeta) + 12ab(U(\zeta)U'(\zeta))' + (-a^3b + ab^3)U^{iv}(\zeta) = 0.$$
 (8)

Integrating Eq.8 twice and neglecting constants of integration

$$(4av - 6cd)U(\zeta) + 6abU^{2}(\zeta) + (-a^{3}b + ab^{3})U''(\zeta) = 0.$$
(9)

Balancing between the highest nonlinear term and the highest derivative U'' with U^2 in Eq. 9 gives N=2. Thus, Eq. 9 has the formal solution

$$U(\zeta) = \eta_0 + \eta_1 P(\zeta) + \eta_2 P^2(\zeta), \qquad (10)$$

(2)



$$\begin{cases} 6ab\,\eta_{2}^{2} + 6ab^{3}\,\eta_{2}\ln^{2}A\Omega^{2} - 6a^{3}b\eta_{2}\ln^{2}A\Omega^{2} = 0, \\ -10\,a^{3}b\,\eta_{2}\ln^{2}A\Theta\,\Omega + 10\,ab^{3}\,\eta_{2}\ln^{2}A\Theta\,\Omega - 2\,a^{3}b\eta_{1}\ln^{2}A\Omega^{2} + 12\,ab\,\eta_{1}\eta_{2} + 2\,ab^{3}\eta_{1}\ln^{2}A\Omega^{2} = 0, \\ 8\,ab^{3}\,\eta_{2}\ln^{2}A\Psi\Omega + 6\,ab\,\eta_{1}^{2} + 12\,ab\,\eta_{0}\eta_{2} + 4\,ab^{3}\eta_{2}\ln^{2}A\Theta^{2} - 8\,a^{3}b\eta_{2}\ln^{2}A\Psi\Omega - 6\,cd\,\eta_{2} + 4\,av\eta_{2} \\ + 3\,ab^{3}\,\eta_{1}\ln^{2}A\Theta\,\Omega - 4\,a^{3}b\eta_{2}\ln^{2}A\Theta^{2} - 3\,a^{3}b\eta_{1}\ln^{2}A\Theta\,\Omega = 0, \\ -6\,a^{3}b\eta_{2}\ln^{2}A\Psi\Theta - 2\,a^{3}b\eta_{1}\ln^{2}A\Omega\,\Psi + 4\,av\eta_{1} - a^{3}b\eta_{1}\ln^{2}A\Theta^{2} - 6\,cd\,\eta_{1} + 6\,ab^{3}\eta_{2}\ln^{2}A\Psi\Theta \\ + 2\,ab^{3}\,\eta_{1}\ln^{2}A\Omega\,\Psi + 12\,ab\,\eta_{0}\eta_{1} + ab^{3}\eta_{1}\ln^{2}A\Theta^{2} = 0, \\ 4\,av\,\eta_{0} + 2\,ab^{3}\,\eta_{2}\ln^{2}A\Psi^{2} + 6\,ab\,\eta_{0}^{2} - a^{3}b\eta_{1}\ln^{2}A\Theta\,\Psi - 6\,cd\,\eta_{0} - 2\,a^{3}b\eta_{2}\ln^{2}A\Psi^{2} \\ + ab^{3}\,\eta_{1}\ln^{2}A\Theta\,\Psi = 0. \end{cases}$$

$$(11)$$

Solving the above system of algebraic Eq. 11 with the aid of Maple, we have the following results;

Set 1

$$\begin{cases} \eta_{0} = \Psi \Omega \left(a^{2} - b^{2} \right) \ln^{2} A, \\ \eta_{1} = \Theta \Omega \left(a^{2} - b^{2} \right) \ln^{2} A, \\ \eta_{2} = \Omega^{2} \left(a^{2} - b^{2} \right) \ln^{2} A, \\ v = \frac{1}{4} b \left(b^{2} - a^{2} \right) \left(\Theta^{2} - 4 \Psi \Omega \right) \ln^{2} A - \frac{3cd}{2a}. \end{cases}$$
(12)

Set 2

$$\begin{cases} \eta_{0} = \frac{1}{6} (a^{2} - b^{2}) (\Theta^{2} + 2\Psi\Omega) \ln^{2}A, \\ \eta_{1} = \Theta \Omega (a^{2} - b^{2}) \ln^{2}A, \\ \eta_{2} = \Omega^{2} (a^{2} - b^{2}) \ln^{2}A, \\ v = \frac{1}{4} b (b^{2} - a^{2}) (\Theta^{2} - 4\Psi\Omega) \ln^{2}A + \frac{3cd}{2a}. \end{cases}$$
(13)

substituting Eq. 12 along with Eq. 10 into Eq. 7, we get the following exact solutions of Eq. 6

Case 1

Family 1. When $\Theta^2 - 4\Psi\Omega < 0$ and $\Omega \neq 0$, then the traveling wave solutions are given by

$$u_{1,1}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \tan_A \left(\frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right] + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \tan_A \left(\frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right]^2$$
(14)

$$u_{2,1}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left[-\frac{\Theta}{2\Omega} - \frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \cot_A \left(\frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right] + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left[-\frac{\Theta}{2\Omega} - \frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \cot_A \left(\frac{\sqrt{-(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right]^2$$
(15)

$$u_{3,1}(\zeta) = \Psi\Omega\left(a^{2} - b^{2}\right)\ln^{2}A + \Theta\Omega\left(a^{2} - b^{2}\right)\ln^{2}A$$

$$\left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{2\Omega}\left(\tan_{A}\left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \pm \sqrt{pq}\sec_{A}\left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)}\zeta\right)\right)\right]$$

$$+ \Omega^{2}\left(a^{2} - b^{2}\right)\ln^{2}A\left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{2\Omega}\left(\tan_{A}\left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \pm \sqrt{pq}\sec_{A}\left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)}\zeta\right)\right)\right]^{2}$$
(16)



$$u_{4,1}(\zeta) = \Psi \Omega \left(a^{2} - b^{2} \right) \ln^{2}A + \Theta \Omega \left(a^{2} - b^{2} \right) \ln^{2}A$$

$$\left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{2\Omega} \left(-\cot_{A} \left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \right) \right]$$

$$+ \Omega^{2} \left(a^{2} - b^{2} \right) \ln^{2}A \left[-\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{2\Omega} \left(-\cot_{A} \left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(\sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \right) \right]^{2}$$

$$(17)$$

$$u_{5,1}(\zeta) = \Psi \Omega \left(a^{2} - b^{2} \right) \ln^{2}A + \Theta \Omega \left(a^{2} - b^{2} \right) \ln^{2}A$$

$$\left\{ -\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{4\Omega} \left(\tan_{A} \left(\frac{1}{4} \sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) - \cot_{A} \left(\frac{1}{4} \sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \right) \right\}$$

$$+ \Omega^{2} \left(a^{2} - b^{2} \right) \ln^{2}A$$

$$\left\{ -\frac{\Theta}{2\Omega} + \frac{\sqrt{-(\Theta^{2} - 4\Psi\Omega)}}{4\Omega} \left(\tan_{A} \left(\frac{1}{4} \sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) - \cot_{A} \left(\frac{1}{4} \sqrt{-(\Theta^{2} - 4\Psi\Omega)} \zeta \right) \right) \right\}^{2}.$$

$$(18)$$

Family2. When $\Theta^2 - 4\Psi\Omega > 0$ and $\Omega \neq 0$, then the traveling wave solutions are given by

$$u_{6,2}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \tanh_A \left(\frac{\sqrt{(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right\} + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \tanh_A \left(\frac{\sqrt{(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right\}^2, \quad (19)$$

$$u_{7,2}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \coth_A \left(\frac{\sqrt{(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right\} + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^2 - 4\Psi\Omega)}}{2\Omega} \coth_A \left(\frac{\sqrt{(\Theta^2 - 4\Psi\Omega)} \zeta}{2} \right) \right\}^2, \quad (20)$$

$$\begin{split} u_{8,2}(\zeta) &= \Psi \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A + \Theta \, \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A \\ & \left\{ -\frac{\Theta}{2\,\Omega} + \frac{\sqrt{(\Theta^2 - 4\Psi\,\Omega)}}{2\,\Omega} \left(-\tanh_A \left(\sqrt{(\Theta^2 - 4\Psi\,\Omega)} \, \zeta \right) \pm i\,\sqrt{p\,q} \quad csch \quad _A \left(\sqrt{(\Theta^2 - 4\Psi\,\Omega)} \, \zeta \right) \right) \right\} + \\ & \Omega^2 \, \left(a^2 - b^2\right) \ln^2\!\! A \\ & \left\{ -\frac{\Theta}{2\,\Omega} + \frac{\sqrt{(\Theta^2 - 4\Psi\,\Omega)}}{2\,\Omega} \left(-\tanh_A \left(\sqrt{(\Theta^2 - 4\Psi\,\Omega)} \, \zeta \right) \pm i\,\sqrt{p\,q} \quad csch \quad _A \left(\sqrt{(\Theta^2 - 4\Psi\,\Omega)} \, \zeta \right) \right) \right\}^2 \end{split} \tag{21}$$

$$u_{9,2}(\zeta) = \Psi\Omega \left(a^{2} - b^{2}\right) \ln^{2}A + \Theta\Omega \left(a^{2} - b^{2}\right) \ln^{2}A$$

$$\left\{ -\frac{\Theta}{2\Omega} + \frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)}}{2\Omega} \left(-\coth_{A}\left(\sqrt{(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \pm \sqrt{pq} \quad csch \quad _{A}\left(\sqrt{(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \right) \right\}$$

$$+ \Omega^{2} \left(a^{2} - b^{2}\right) \ln^{2}A$$

$$\left\{ -\frac{\Theta}{2\Omega} + \frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)}}{2\Omega} \left(-\coth_{A}\left(\sqrt{(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \pm \sqrt{pq} \quad csch \quad _{A}\left(\sqrt{(\Theta^{2} - 4\Psi\Omega)}\zeta\right) \right) \right\}^{2}$$
 (22)

$$u_{10,2}(\zeta) = \Psi \Omega \left(a^{2} - b^{2}\right) \ln^{2}A + \Theta \Omega \left(a^{2} - b^{2}\right) \ln^{2}A$$

$$\left[-\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)}}{4\Omega} \left\{ \tanh_{A} \left(\frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)} \zeta}{4} \right) + \coth_{A} \left(\frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)} \zeta}{4} \right) \right\} \right] + \Omega^{2} \left(a^{2} - b^{2}\right) \ln^{2}A$$

$$\left[-\frac{\Theta}{2\Omega} - \frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)}}{4\Omega} \left\{ \tanh_{A} \left(\frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)} \zeta}{4} \right) + \coth_{A} \left(\frac{\sqrt{(\Theta^{2} - 4\Psi\Omega)} \zeta}{4} \right) \right\} \right]^{2}. \quad (23)$$



Family3. When $\Psi \Omega > 0$ and $\Theta = 0$, then the traveling wave solutions are given by

$$u_{11,3}(\zeta) = \Psi \Omega \left(a^{2} - b^{2}\right) \ln^{2} A + \Theta \Omega \left(a^{2} - b^{2}\right) \ln^{2} A \left\{ \sqrt{\frac{\Psi}{\Omega}} \tan_{A} \left(\sqrt{\Psi \Omega} \zeta\right) \right\} + \Omega^{2} \left(a^{2} - b^{2}\right) \ln^{2} A \left\{ \sqrt{\frac{\Psi}{\Omega}} \tan_{A} \left(\sqrt{\Psi \Omega} \zeta\right) \right\}^{2}, \quad (24)$$

$$u_{12,3}(\zeta) = \Psi \Omega \left(a^{2} - b^{2}\right) \ln^{2} A + \Theta \Omega \left(a^{2} - b^{2}\right) \ln^{2} A \left\{-\sqrt{\frac{\Psi}{\Omega}} \cot_{A}\left(\sqrt{\Psi \Omega}\zeta\right)\right\} + \Omega^{2} \left(a^{2} - b^{2}\right) \ln^{2} A \left\{-\sqrt{\frac{\Psi}{\Omega}} \cot_{A}\left(\sqrt{\Psi \Omega}\zeta\right)\right\}^{2}, \quad (25)$$

$$\begin{split} u_{13,3}\left(\zeta\right) &= \Psi \Omega \left(a^2 - b^2\right) \ln^2\!\! A + \Theta \Omega \left(a^2 - b^2\right) \ln^2\!\! A \left\{ \sqrt{\frac{\Psi}{\Omega}} \left(\tan_A \left(2\sqrt{\Psi \Omega} \zeta \right) \pm \sqrt{pq} \sec_A \left(2\sqrt{\Psi \Omega} \zeta \right) \right) \right\} \\ &+ \Omega^2 \left(a^2 - b^2\right) \ln^2\!\! A \left\{ \sqrt{\frac{\Psi}{\Omega}} \left(\tan_A \left(2\sqrt{\Psi \Omega} \zeta \right) \pm \sqrt{pq} \sec_A \left(2\sqrt{\Psi \Omega} \zeta \right) \right) \right\}^2, \end{split} \tag{26}$$

$$u_{14,3}(\zeta) = \Psi \Omega \left(a^{2} - b^{2} \right) \ln^{2}A + \Theta \Omega \left(a^{2} - b^{2} \right) \ln^{2}A \left\{ \sqrt{\frac{\Psi}{\Omega}} \left(-\cot_{A} \left(2\sqrt{\Psi \Omega} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(2\sqrt{\Psi \Omega} \zeta \right) \right) \right\} + \Omega^{2} \left(a^{2} - b^{2} \right) \ln^{2}A \left\{ \sqrt{\frac{\Psi}{\Omega}} \left(-\cot_{A} \left(2\sqrt{\Psi \Omega} \zeta \right) \pm \sqrt{pq} \csc_{A} \left(2\sqrt{\Psi \Omega} \zeta \right) \right) \right\}^{2}, \quad (27)$$

$$u_{15,3}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ \frac{1}{2} \sqrt{\frac{\Psi}{\Omega}} \left(\tan_A \left(\frac{1}{2} \sqrt{\Psi \Omega} \zeta \right) - \cot_A \left(\frac{1}{2} \sqrt{\Psi \Omega} \zeta \right) \right) \right\}$$

$$+ \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ \frac{1}{2} \sqrt{\frac{\Psi}{\Omega}} \left(\tan_A \left(\frac{1}{2} \sqrt{\Psi \Omega} \zeta \right) - \cot_A \left(\frac{1}{2} \sqrt{\Psi \Omega} \zeta \right) \right) \right\}^2. \quad (28)$$

Family4. When $\Psi\Omega<0$ and $\Theta=0$, then the traveling wave solutions are given by

$$\begin{split} u_{16,4}\left(\zeta\right) &= \Psi \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A + \Theta \, \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A \, \left\{ -\sqrt{-\frac{\Psi}{\Omega}} \tanh_A \left(\sqrt{-\Psi \Omega} \zeta\right) \right\} \\ &+ \Omega^2 \left(a^2 - b^2\right) \ln^2\!\! A \left\{ -\sqrt{-\frac{\Psi}{\Omega}} \tanh_A \left(\sqrt{-\Psi \Omega} \zeta\right) \right\}^2, \quad (29) \end{split}$$

$$u_{17,4}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-\sqrt{-\frac{\Psi}{\Omega}} \coth_A\left(\sqrt{-\Psi \Omega}\zeta\right)\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-\sqrt{-\frac{\Psi}{\Omega}} \coth_A\left(\sqrt{-\Psi \Omega}\zeta\right)\right\}^2, \quad (30)$$

$$\begin{split} u_{18,4}\left(\zeta\right) &= \Psi \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A + \Theta \, \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A \left\{ \sqrt{-\frac{\Psi}{\Omega}} \left(-\tanh \left(2\sqrt{-\Psi\Omega}\zeta\right) \right) \pm i\sqrt{pq} \quad csch \quad {}_A(2\sqrt{-\Psi\Omega}\zeta) \right\} \right. \\ &\left. + \Omega^2 \left(a^2 - b^2\right) \ln^2\!\! A \left\{ \sqrt{\frac{-\Psi}{\Omega}} \left(-\tanh \left(2\sqrt{-\Psi\Omega}\zeta\right) \right) \pm i\sqrt{pq} \quad csch \quad {}_A(2\sqrt{-\Psi\Omega}\zeta) \right\}^2, \quad (31) \end{split}$$



$$\begin{split} u_{19,4}\left(\zeta\right) &= \Psi \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A + \Theta \, \Omega \, \left(a^2 - b^2\right) \ln^2\!\! A \, \left\{ -\sqrt{\frac{\Psi}{\Omega}} \left(-\coth_A \left(2\sqrt{-\Psi\,\Omega}\,\zeta\right) \pm \sqrt{pq} \quad csch \quad _A \left(2\sqrt{-\Psi\,\Omega}\,\zeta\right) \right) \right\} \\ &+ \Omega^2 \left(a^2 - b^2\right) \ln^2\!\! A \left\{ \sqrt{-\frac{\Psi}{\Omega}} \left(-\coth_A \left(2\sqrt{-\Psi\,\Omega}\,\zeta\right) \pm \sqrt{pq} \quad csch \quad _A \left(2\sqrt{-\Psi\,\Omega}\,\zeta\right) \right) \right\}^2 \end{split} \tag{32}$$

$$u_{20,4}(\zeta) = \Psi \Omega \left(a^{2} - b^{2}\right) \ln^{2}A + \Theta \Omega \left(a^{2} - b^{2}\right) \ln^{2}A \left\{-\frac{1}{2} \sqrt{-\frac{\Psi}{\Omega}} \left(\tanh_{A} \left(\frac{1}{2} \sqrt{-\Psi \Omega} \zeta\right) + \coth_{A} \sqrt{-\Psi \Omega} \zeta\right)\right\} + \Omega^{2} \left(a^{2} - b^{2}\right) \ln^{2}A \left\{-\frac{1}{2} \sqrt{-\frac{\Psi}{\Omega}} \left(\tanh_{A} \left(\frac{1}{2} \sqrt{-\Psi \Omega} \zeta\right) + \coth_{A} \sqrt{-\Psi \Omega} \zeta\right)\right\}^{2}\right\}.$$
(33)

Family5. When $\Theta = 0$ and $\Omega = \Psi$, then the traveling wave solutions are given by

$$u_{21,5}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ \tan_A (\Psi \zeta) \right\} + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ \tan_A (\Psi \zeta) \right\}^2, \tag{34}$$

$$u_{22,5}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ -\cot_A (\Psi \zeta) \right\} + \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ -\cot_A (\Psi \zeta) \right\}^2, \tag{35}$$

$$u_{23,5}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{ \left(\tan_A \left(2\Psi\zeta\right) \pm \sqrt{pq}\right) \sec_A \left(2\Psi\zeta\right) \right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{ \left(\tan_A \left(2\Psi\zeta\right) \pm \sqrt{pq}\right) \sec_A \left(2\Psi\zeta\right) \right\}^2, \quad (36)$$

$$u_{24,5}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-\cot_A \left(2\Psi\zeta\right) \pm \sqrt{pq} \csc_A \left(2\Psi\zeta\right)\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-\cot_A \left(2\Psi\zeta\right) \pm \sqrt{pq} \csc_A \left(2\Psi\zeta\right)\right\}^2, \quad (37)$$

$$u_{25,5}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{ \frac{1}{2} \left(\tan \left(\frac{1}{2} \Psi \zeta \right) - \cot \left(\frac{1}{2} \Psi \zeta \right) \right) \right\}$$

$$+ \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{ \frac{1}{2} \left(\tan \left(\frac{1}{2} \Psi \zeta \right) - \cot \left(\frac{1}{2} \Psi \zeta \right) \right) \right\}^2. \quad (38)$$

Family6. When $\Theta = 0$ and $\Omega = -\Psi$, then the traveling wave solutions are given by

$$u_{26,6}\left(\zeta\right) = \Psi\Omega\left(a^2 - b^2\right)\ln^2\!A + \Theta\Omega\left(a^2 - b^2\right)\ln^2\!A\left\{-\tanh_A\left(\Psi\zeta\right)\right\} + \Omega^2\left(a^2 - b^2\right)\ln^2\!A\left\{-\tanh_A\left(\Psi\zeta\right)\right\}^2, \tag{39}$$

$$u_{27,6}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-\coth_A \left(\Psi \zeta\right)\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-\coth_A \left(\Psi \zeta\right)\right\}^2,\tag{40}$$

$$u_{28,6}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-\tanh_A \left(2\Psi \zeta\right) \pm i\sqrt{pq} \operatorname{sech}_A \left(2\Psi \zeta\right)\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-\tanh_A \left(2\Psi \zeta\right) \pm i\sqrt{pq} \operatorname{sech}_A \left(2\Psi \zeta\right)\right\}^2, \quad (41)$$

$$u_{29,6}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-\coth_A \left(2\Psi\zeta\right) \pm \sqrt{pq} \operatorname{csch}_A \left(2\Psi\zeta\right)\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-\coth_A \left(2\Psi\zeta\right) \pm \sqrt{pq} \operatorname{csch}_A \left(2\Psi\zeta\right)\right\}^2, \quad (42)$$



$$u_{30,6}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{1}{2} \left(\tanh_A \left(\frac{1}{2} \Psi \zeta \right) + \coth_A \left(\frac{1}{2} \Psi \zeta \right) \right) \right\}$$

$$+ \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ -\frac{1}{2} \left(\tanh_A \left(\frac{1}{2} \Psi \zeta \right) + \coth_A \left(\frac{1}{2} \Psi \zeta \right) \right) \right\}^2. \tag{43}$$

Family7 When $\Theta^2 = 4\Psi \Omega$, then the traveling wave solutions are given by

$$u_{31,7}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{-2 \frac{\Psi(\Theta \zeta \ln A + 2)}{\Theta^2 \zeta \ln A}\right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{-2 \frac{\Psi(\Theta \zeta \ln A + 2)}{\Theta^2 \zeta \ln A}\right\}^2$$
(44)

Family 8. When $\Theta = 0, \Psi = mk, (m \neq 0)$ and $\Omega = k$, then the traveling wave solutions are given by

$$u_{32,8}(\zeta) = \Psi \Omega \left(a^2 - b^2 \right) \ln^2 A + \Theta \Omega \left(a^2 - b^2 \right) \ln^2 A \left\{ A^{k\zeta} - m \right\}$$

$$+ \Omega^2 \left(a^2 - b^2 \right) \ln^2 A \left\{ A^{k\zeta} - m \right\}^2$$
(45)

Family 9. When $\Theta = \Omega = 0$,, then the traveling wave solutions are given by

$$u_{33,9}\left(\zeta\right) = \Psi\Omega\left(a^2 - b^2\right)\ln^2 A + \Theta\Omega\left(a^2 - b^2\right)\ln^2 A\left\{\Psi\zeta\ln(A)\right\} + \Omega^2\left(a^2 - b^2\right)\ln^2 A\left\{\Psi\zeta\ln(A)\right\}^2 \tag{46}$$

Family 10: when $\Theta = \Psi = 0$,, then

$$u_{34,10}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{\frac{-1}{\Omega \zeta \ln(A)}\right\}$$

$$+ \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{\frac{-1}{\Omega \zeta \ln(A)}\right\}^2$$

$$(47)$$

Family 11: When $\Theta \neq 0, \Psi = 0$, then the traveling wave solutions are given by

$$u_{35,11}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{p \Theta}{\Omega \left(\cosh_A(\Theta \zeta) - \sinh_A(\Theta \zeta)\right) + p} \right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{p \Theta}{\Omega \left(\cosh_A(\Theta \zeta) - \sinh_A(\Theta \zeta) + p\right)} \right\}^2, \quad (48)$$

$$u_{36,11}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{\Theta \left(\sinh_A(\Theta \zeta) + \cosh_A(\Theta \zeta)\right)}{\Omega \left(\sinh_A(\Theta \zeta) + \cosh_A(\Theta \zeta) + q\right)} \right\} \\ + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{\Theta \left(\sinh_A(\Theta \zeta) + \cosh_A(\Theta \zeta)\right)}{\Omega \left(\sinh_A(\Theta \zeta) + \cosh_A(\Theta \zeta) + q\right)} \right\}^2. \tag{49}$$

Family 12: When $\Theta = \lambda$, $\Omega = m (m \neq 0)$, and $\Psi = 0$ then the rational solution is given by

$$u_{37,12}(\zeta) = \Psi \Omega \left(a^2 - b^2\right) \ln^2 A + \Theta \Omega \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{pA^{\lambda \zeta}}{q - mpA^{\lambda \zeta}} \right\} + \Omega^2 \left(a^2 - b^2\right) \ln^2 A \left\{ -\frac{pA^{\lambda \zeta}}{q - mpA^{\lambda \zeta}} \right\}^2$$
(50)

where ζ is an independent variable, p and q are arbitrary constants greater than zero and called deformation parameters.

4 The graphical representation






Fig. 1: The 3-D and 2-D graphs of $u_1(\zeta)$ given by Eq. 14.




Fig. 2: The 3-D and 2-D graphs of $u_2(\zeta)$ given by Eq. 15.




Fig. 3: The 3-D and 2-D graphs of $u_3(\zeta)$ given by Eq.16.






Fig. 4: The 3-D and 2-D graphs of $u_4(\zeta)$ given by Eq. 17.




Fig. 5: The 3-D and 2-D graphs of $u_5(\zeta)$ given by Eq. 18.




Fig. 6: The 3-D and 2-D graphs of $u_6(\zeta)$ given by Eq. 19.




Fig. 7: The 3-D and 2-D graphs of $u_7(\zeta)$ given by Eq. 20.




Fig. 8: The 3-D and 2-D graphs of $u_8(\zeta)$ given by Eq. 21.




Fig. 9: The 3-D and 2-D graphs of $u_9(\zeta)$ given by Eq. 22.






Fig. 10: The 3-D and 2-D graphs of $u_{10}(\zeta)$ given by Eq. 23.




Fig. 11: The 3-D and 2-D graphs of $u_{11}(\zeta)$ given by Eq. 24.




Fig. 12: The 3-D and 2-D graphs of $u_{12}(\zeta)$ given by Eq. 25.




Fig. 13: The 3-D and 2-D graphs of $u_{13}(\zeta)$ given by Eq. 26.




Fig. 14: The 3-D and 2-D graphs of $u_{15}(\zeta)$ given by Eq. 28.




Fig. 15: The 3-D and 2-D graphs of $u_{16}(\zeta)$ given by Eq. 29.






Fig. 16: The 3-D and 2-D graphs of $u_{17}(\zeta)$ given by Eq. 30.




Fig. 17: The 3-D and 2-D graphs of $u_{19}(\zeta)$ given by Eq. 32.




Fig. 18: The 3-D and 2-D graphs of $u_{20}(\zeta)$ given by Eq. 33.






Fig. 19: The 3-D and 2-D graphs of $u_{21}(\zeta)$ given by Eq. 34.




Fig. 20: The 3-D and 2-D graphs of $u_{22}(\zeta)$ given by Eq. 35.




Fig. 21: The 3-D and 2-D graphs of $u_{23}(\zeta)$ given by Eq. 36.






Fig. 22: The 3-D and 2-D graphs of $u_{24}(\zeta)$ given by Eq. 37.




Fig. 23: The 3-D and 2-D graphs of $u_{25}(\zeta)$ given by Eq. 38.




Fig. 24: The 3-D and 2-D graphs of $u_{26}(\zeta)$ given by Eq. 39.






Fig. 25: The 3-D and 2-D graphs of $u_{27}(\zeta)$ given by Eq. 40.




Fig. 26: The 3-D and 2-D graphs of $u_{29}(\zeta)$ given by Eq. 42.




Fig. 27: The 3-D and 2-D graphs of $u_{31}(\zeta)$ given by Eq. 44.






Fig. 28: The 3-D and 2-D graphs of $u_{32}(\zeta)$ given by Eq. 45.




Fig. 29: The 3-D and 2-D graphs of $u_{34}(\zeta)$ given by Eq. 47.



5 Physical Interpretation

equation The (4+1)-dimensional Fokas is higher-dimensional extension of the well-known Kadomtsev-Petviashvili equation, models which nonlinear wave propagation in dispersive media. It finds applications in various fields, including fluid mechanics, plasma physics, and nonlinear optics. The traveling wave solutions obtained in this study represent different types of wave patterns that can emerge in the systems governed by this equation. Soliton solutions, for example, are associated with temporary, real, localized wave patterns which do not deform or change their velocity while propagating. Periodic solutions are those of wave like form with some form of repetition whereas rational solutions correspond to localized disturbances which decay algebraically. Such solutions are highly useful for understanding the behavior of physical systems that underlie individual waves and can be applied for the computation of many wave processes.

6 Discussion and Conclusion

In the present study, we were able to implement the recent proposed new extended direct algebraic method to obtain a general solution for the traveling wave solutions of the (4 + 1)-dimensional Fokas equation. The acquires solutions contain solitons, periodical functions, rational function and so on. In doing so, the current work makes a contribution to knowledge of the various solutions of this largely nonlinear partial differential equation and lest a potential application in the corresponding fields. The effectiveness of the proposed method is shown in terms of solving high-dimensional nonlinear PDEs while it can be used for other nonlinear systems as well. further investigations can be the stability analysis of the solutions, the physical interpretation of the solutions and last but not the least, numerical studies for the validation of the theory.

References

- [1] STR. Rizvi, AR. Seadawy, F. Ashraf, M. Younis, H. Iqbal, and D. Baleanu, 2020. Lump and interaction solutions of a geophysical Korteweg-de Vries equation. Results in Physics, 19: 103661.
- [2] Farrah Ashraf, Aly R.Seadawy, Syed T.R.Rizvi, Kashif Ali, and M. Aamir Ashraf, 2022. Multi-wave, M-shaped rational and interaction solutions for fractional nonlinear electrical transmission line equation. Journal of Geometry and Physics, 177: 104503.
- [3] Rizvi. STR, Seadawy. AR, Ashraf. F, Younis .M, Iqbal. H, and Baleanu. D, 2019. Kinky breathers, W-shaped and multi-peak solitons interaction in (2 + 1)-dimensional nonlinear Schrodinger equation with Kerr law of nonlinearity. Eur. Phys. J. Plus, 134: 1-10.

- [4] Romana Ashraf, Faiza Amanat, Farah Ashraf, Soliton solutions for the (4+1)-dimensional Fokas equation using integration techniques, Alexandria Engineering Journal, 107 (2024) 61-72.
- [5] Ali Akgül, Saliha Manzoor, Farrah Ashraf, Romana Ashraf, Exact solutions of the (2+1)-dimensional Zoomeron model arising in nonlinear optics via mapping method, Optical and Quantum Electronics, 56 (7) (2024) 1207.
- [6] Shabbir Hussain, Muhammad Sajid Iqbal, Mustafa Bayram, Romana Ashraf, Mustafa Inc, Shahram Rezapour, Muhammad Akhtar Tarar, Optical soliton solutions in a distinctive class of nonlinear Schrödinger's equation with cubic, quintic, septic, and nonic nonlinearities, Optical and Quantum Electronics, 56 (6) (2024) 1066.
- [7] Mustafa Inc, Shabbir Hussain, Ali Hasan Ali, Muhammad Sajid Iqbal, Romana Ashraf, Muhammad Akhtar Tarar, Muhammad Adnan, Analyzing solitary wave solutions of the nonlinear Murray equation for blood flow in vessels with non-uniform wall properties, Scientific Reports, 10588 (2024).
- [8] F. Ashraf, R. Ashraf, A. Akgül, Travelling waves solutions of Hirota-Ramani equation by modified extended direct algebraic method and new extended direct algebraic method, International Journal of Modern Physics B, (2024).
- [9] R. Ashraf, S. Hussain, M. S. Iqbal, The extended Fan's sub-equation method and its application to nonlinear Schrödinger equation with saturable nonlinearity, Results in Physics, (2023).
- [10] S. Hussain, M. S. Iqbal, R. Ashraf, M. Inc, M. A. Tarar, Exploring nonlinear dispersive waves in a disordered medium: an analysis using ?6 model expansion method, Optical and Quantum Electronics, 55 (7) (2023) 651.
- [11] S. Hussain, M. S. Iqbal, R. Ashraf, M. Inc, M. A. Tarar, Quantum analysis of nonlinear optics in Kerr affected saturable nonlinear media and multiplicative noise: a path to new discoveries, Optical and Quantum Electronics, 55 (7) (2023) 578.
- [12] R. Ashraf, F. Ashraf, A Akgül, S Ashraf, B Alshahrani, M Mahmoud, Some new soliton solutions to the (3 + 1)-dimensional generalized KdV-ZK equation via enhanced modified extended tanh-expansion approach, Alexandria Engineering Journal, 69 (2023) 303-309.
- [13] S.T. R. Rizvi, Aly R. Seadawy, Sarfaraz Ahmed, R. Ashraf, Lax pair, Darboux transformation, Weierstrass-Jacobi elliptic and generalized breathers along with soliton solutions for Benjamin Bona Mahony equation, International Journal of Modern Physics B, (2023) 2350233.
- [14] Aly R. Seadawy, S.T. R. Rizvi, Tahira Batool, R. Ashraf, Study of Sasa-Satsuma dynamical system for Kuznetsov-Ma and generalized breathers, lump, periodic and rogue wave solutions, International Journal of Modern Physics B, (2023) 2350181.
- [15] Farrah Ashraf, Tehsina Javeed, Romana Ashraf, Amina Rana, Ali Akgul, Some new soliton solution to the higher dimensional burger-huxley and shallow water waves equation with couple of integration architectonic, Results in Physics, 43 (2022) 106048.
- [16] S.T. R. Rizvi, Aly R. Seadawy, R. Ashraf, Propagation of chirped periodic and solitary waves for the coupled nonlinear Schrödinger equation in two core optical fibers



- with parabolic law with weak non-local nonlinearity, Optical and Quantum Electronics, 545 (2022).
- [17] S. Ahmed, R. Ashraf, Aly R. Seadawy, S.T. R. Rizvi, M. Younis, Ali Althobaiti, Ahmed M. El-Shehawi, Lump, multi-wave, Kinky breathers, interactional solutions, and stability analysis for general (2+1)-th dispersionless Dym equation, Results in Physics, 25 (2021) 104160.
- [18] R. Ashraf, M.O. Ahmad, M. Younis, K.U. Tariq, K. Ali, S.T.R. Rizvi: Dipole and combo solitons in DWDM systems, Optik, 158 (2018) 1073-1079.
- [19] R. Ashraf, M.O. Ahmad, M. Younis, K. Ali, S.T.R. Rizvi: Dipole and Gausson soliton for ultrashort laser pulse with high order dispersion, Superlattices and Microstructures, 109 (2017) 504-510.
- [20] S.T.R. Rizvi, S. Bashir, M. Younis, R. Ashraf, M.O. Ahmad: Ahmad: Exact soliton of (2 + 1)-dimensional fractional Schrödinger equation, Superlattices and Microstructures, 107 (2017) 234-239.
- [21] Alkhidhr, H. A., & Abdelrahman, M. A. (2021). Wave structures to the three coupled nonlinear Maccaris systems in plasma physics. Results in Physics, 105092.
- [22] Ziane, D., Hamdi Cherif, M., Baleanu, D., & Belghaba, K. (2020). Non-Differentiable Solution of Nonlinear Biological Population Model on Cantor Sets. Fractal and Fractional, 4(1), 5.
- [23] Younis, M., Sulaiman, T. A., Bilal, M., Rehman, S. U., & Younas, U. (2020). Modulation instability analysis, optical and other solutions to the modified nonlinear Schrodinger equation. Communications in Theoretical Physics, 72(6), 065001.
- [24] Younas, U., Seadawy, A. R., Younis, M., & Rizvi, S. T. R. (2020). Dispersive of propagation wave structures to the Dullin-Gottwald-Holm dynamical equation in a shallow water waves. Chinese Journal of Physics, 68, 348-364.
- [25] Kim, H., Bae, J. H., & Sakthivel, R. (2014). Exact travelling wave solutions of two important nonlinear partial differential equations. Zeitschrift fur Naturforschung A, 69(3-4), 155-162.
- [26] Paredes, A., Olivieri, D. N., & Michinel, H. (2020). From optics to dark matter: A review on nonlinear Schrodinger-Poisson systems. Physica D: Nonlinear Phenomena, 403, 132301.
- [27] Yomba, E. (2005). The extended Fan's sub-equation method and its application to KdV-MKdV, BKK and variant Boussinesq equations. Physics Letters A, 336(6), 463-476.
- [28] Batool, F., & Akram, G. (2017). Application of extended Fan sub-equation method to (1+ 1)-dimensional nonlinear dispersive modified Benjamin-Bona-Mahony equation with fractional evolution. Optical and Quantum Electronics, 49(11), 1-9.
- [29] El-Wakil, S. A., & Abdou, M. A. (2008). The extended Fan sub-equation method and its applications for a class of nonlinear evolution equations. Chaos, Solitons & Fractals, 36(2), 343-353.
- [30] Cheemaa, N., & Younis, M. (2016). New and more exact traveling wave solutions to integrable (2+ 1)-dimensional Maccari system. Nonlinear Dynamics, 83(3), 1395-1401.
- [31] Lakkis, O., & Pryer, T. (2013). A finite element method for nonlinear elliptic problems. SIAM Journal on Scientific Computing, 35(4), A2025-A2045.

- [32] Yan, S., & Jin, J. M. (2015). Theoretical formulation of a time-domain finite element method for nonlinear magnetic problems in three dimensions. Progress In Electromagnetics Research, 153, 33-55.
- [33] Carrillo, J. A., Chertock, A., & Huang, Y. (2015). A finite-volume method for nonlinear nonlocal equations with a gradient flow structure. Communications in Computational Physics, 17(1), 233-258.
- [34] Murakawa, H. (2017). A linear finite volume method for nonlinear cross-diffusion systems. Numerische Mathematik, 136(1), 1-26.
- [35] Gavete, L., Urea, F., Benito, J. J., Garca, A., Urea, M., & Salete, E. (2017). Solving second order non-linear elliptic partial differential equations using generalized finite difference method. Journal of Computational and Applied Mathematics, 318, 378-387.
- [36] Urea, F., Gavete, L., Garcia, A., Benito, J. J., & Vargas, A. M. (2019). Solving second order non-linear parabolic PDEs using generalized finite difference method (GFDM). Journal of Computational and Applied Mathematics, 354, 221-241.
- [37] Oru. (2020). An efficient wavelet collocation method for nonlinear two-space dimensional Fisher-Kolmogorov-Petrovsky-Piscounov equation and two-space dimensional extended Fisher-Kolmogorov equation. Engineering with Computers, 36(3), 839-856.
- [38] Wu, H., Wang, Y., & Zhang, W. (2018). Numerical solution of a class of nonlinear partial differential equations by using barycentric interpolation collocation method. Mathematical Problems in Engineering, 2018.
- [39] Ucar, Y., Karaagac, B., & Esen, A. (2015). A new approach on numerical solutions of the Improved Boussinesq type equation using quadratic B-spline Galerkin finite element method. Applied Mathematics and Computation, 270, 148-155.
- [40] Li, M., Huang, C., & Wang, P. (2017). Galerkin finite element method for nonlinear fractional Schrodinger equations. Numerical Algorithms, 74(2), 499-525.
- [41] Abdusalam, H. A. (2005). On an improved complex tanh-function method. International Journal of Nonlinear Sciences and Numerical Simulation, 6(2), 99-106.
- [42] Aminikhah, H., Moosaei, H., & Hajipour, M. (2010). Exact solutions for nonlinear partial differential equations via Expfunction method. Numerical Methods for Partial Differential Equations, 26(6), 1427-1433.
- [43] Zhao, Y. M. (2013). F-expansion method and its application for finding new exact solutions to the Kudryashov-Sinelshchikov equation. Journal of Applied Mathematics, 2013
- [44] Wazwaz, A. M. (2008). The Hirota s direct method for multiple-soliton solutions for three model equations of shallow water waves. Applied Mathematics and Computation, 201(1-2), 489-503.
- [45] Ryabov, P. N., Sinelshchikov, D. I., & Kochanov, M. B. (2011). Application of the Kudryashov method for finding exact solutions of the high order nonlinear evolution equations. Applied Mathematics and Computation, 218(7), 3965-3972.
- [46] Younis, M., Seadawy, A. R., Baber, M. Z., Yasin, M. W., Rizvi, S. T., & Iqbal, M. S. Abundant solitary wave structures of the higher dimensional Sakovich dynamical model. Mathematical Methods in the Applied Sciences.



- [47] Jiong, S. (2003). Auxiliary equation method for solving nonlinear partial differential equations. Physics Letters A, 309(5-6), 387-396.
- [48] Vitanov, N. K. (2011). On modified method of simplest equation for obtaining exact and approximate solutions of nonlinear PDEs: the role of the simplest equation. Communications in Nonlinear Science and Numerical Simulation, 16(11), 4215-4231.
- [49] Mirhosseini-Alizamini, S., Rezazadeh, H., Srinivasa, K. & Bekir, A. New closed form solutions of the new coupled Konnoâ Oono equation using the new extended direct algebraic method. Pramana. 94, 1-12 (2020)
- [50] Rezazadeh, H., Tariq, H., Eslami, M., Mirzazadeh, M. & Zhou, Q. New exact solutions of nonlinear conformable time-fractional Phi-4 equation. Chinese Journal Of Physics. **56**, 2805-2816 (2018)
- [51] Khan, A., Seadawy, A. R., & Nadeem, M. (2021). Soliton solutions for the fractional complex Ginzburg-Landau equation using the Kudryashov method. Applied Mathematics & Information Sciences, 15(1), 73-79.
- [52] Nisar, K. S., Kilicman, A., & Agarwal, P. (2018). Optical soliton solutions for the perturbed Gerdjikov-Ivanov equation with Kerr law nonlinearity. Applied Mathematics & Information Sciences, 12(6), 1233-1240.
- [53] Baleanu, D., Jajarmi, A., & Asad, J. H. (2018). New aspects of the fractional optimal control problems: Mittag-Leffler stability and exact solutions. Progress in Fractional Differentiation and Applications, 4(3), 161-172.
- [54] Gómez-Aguilar, J. F., & Atangana, A. (2017). Fractional Schrödinger equation with non-local and non-singular kernel: A model for nonlinear waves in complex media. Progress in Fractional Differentiation and Applications, 3(2), 85-98.
- [55] El-Depsy, A., & Al-Marzoug, S. M. (2021). Optical Solitons in Nonlinear Thin-Film Waveguides: A Theoretical Approach. International Journal of Thin Film Science and Technology, 10(2), 45-53.
- [56] Rahman, M. M., & Hossain, M. A. (2020). Mathematical Modeling of Heat Transfer in Nanoscale Thin Films Using Fractional PDEs. International Journal of Thin Film Science and Technology, 9(3), 87-95.
- [57] Rahman, M., & Basak, S. (2022). Statistical Properties of Soliton Solutions in the (3+1)-Dimensional Nonlinear Schrödinger Equation. Journal of Statistics Applications & Probability, 11(1), 45-58.
- [58] Alzaatreh, A., & Al-Labadi, L. (2021). A Statistical Approach to Modeling Nonlinear Wave Phenomena with Applications to Soliton Solutions. Journal of Statistics Applications & Probability, 10(2), 215-225.
- [59] Iqbal, M. S. (2011). Solutions of boundary value problems for nonlinear partial differential equations by fixed point methods, T.U. Graz library.
- [60] Rehman, H., Seadawy, A., Younis, M., Rizvi, S., Anwar, I., Baber, M. & Althobaiti, A. Weakly nonlinear electronacoustic waves in the fluid ions propagated via a (3+1)dimensional generalized Kortewegâ de-Vriesâ Zakharovâ Kuznetsov equation in plasma physics. Results In Physics. **33** pp. 105069 (2022)
- [61] Yakada, S., Depelair, B., Betchewe, G. & Doka, S. Miscellaneous new traveling waves in metamaterials by means of the new extended direct algebraic method. Optik. 197 pp. 163108 (2019)

- [62] Mirhosseini-Alizamini, S., Rezazadeh, H., Eslami, M., Mirzazadeh, M. & Korkmaz, A. New extended direct algebraic method for the Tzitzica type evolution equations arising in nonlinear optics. Computational Methods For Differential Equations. 8, 28-53 (2020)
- [63] Vahidi, J., Zabihi, A., Rezazadeh, H. & Ansari, R. New extended direct algebraic method for the resonant nonlinear SchrĶdinger equation with Kerr law nonlinearity. Optik. **227** pp. 165936 (2021)
- [64] Younis, M., Seadawy, A., Baber, M., Husain, S., Iqbal, M., Rizvi, S. & Baleanu, D. Analytical optical soliton solutions of the Schrodinger-Poisson dynamical system. Results In Physics. 27 pp. 104369 (2021)
- [65] Kaplan, M., Akbulut, A. Raza, N. Research on sensitivity analysis and traveling wave solutions of the (4+1)dimensional nonlinear Fokas equation via three different techniques. Physica Scripta. 97, 015203 (2022)



Ahmed Mohamed Hafez earned his Ph. D. degree in Civil Engineering from University of Malaya. He is an Assoc. Prof. at the Department Civil Engineering, INTI International University. He has published over 55 papers in journals. His research

interests are focused on Dam risk, slope stability and soft ground improvement.



Romana **Ashraf** an assistant professor The University at Lahore, Department Computer Science & IT. She is also working as Teaching Area InCharge (TAI) in the respective department. Her research areas are Nonlinear Partial Differential Equations,

Soliton Theory, Numerical Methods, Mathematical Modeling, and Fractional Differential Equations. She has published 21 research papers in internationally reputed journals, which have been extensively cited by researchers both nationally and internationally, with over 281 citations. She has a strong research profile, reflected by an h-index of 9 and an i10-index of 9. She has received the Research Productivity Award 2023 by The University of Lahore, Lahore.





Ali Akgül is a full professor in Siirt university, Faculty of Art and Science, Department of Mathematics. He is the head of the Mathematics department. His research areas are Fractional Differential Equations, Numerical Methods, Partial Differential Equations,

Mathematical Modeling and Functional Analysis. He made a big contribution on fractional calculus and numerical methods. He has more than 700 research papers in very good journals. He has given many talks as an invited speaker in many international conferences. He opened many special issues in very good journals. He is among the World's Top 2% Scientists by Stanford University in 2021, 2022, 2023 and 2024. He got OBADA prize in 2022 (Young Distinguished Researchers).



Montasir Qasymeh received the Ph.D. degree in electrical engineering from Dalhousie University, Halifax, Canada, in 2010, and completed a postdoctoral fellowship at the University of Ottawa, Canada, in 2011. He is currently the Associate Provost for Research,

Innovation, and Academic Development at Abu Dhabi University (ADU), United Arab Emirates, where he also serves as a Professor of Electrical Engineering. Dr. Oasymeh has played a transformative role in advancing ADU's research and innovation agenda. He led the establishment of the MENA region's first quantum computing laboratory, two interdisciplinary research institutes, and the Abu Dhabi Graphene Center. He has also developed robust frameworks to support industry-academia collaboration, startup incubation, and intellectual property commercialization. His research interests include quantum systems, nanoplasmonics, and terahertz wave engineering, with a particular focus on using two-dimensional graphene materials for the design of quantum and terahertz devices. He has authored and co-authored more than 70 peer-reviewed publications, including collaborative work on plasmonic sensors and devices. A prolific inventor, Dr. Qasymeh holds 17 U.S. patents and serves on the editorial board of a leading multidisciplinary journal. He has organized and chaired several high-impact conferences, including serving as General Co-Chair of the International Conference on Electrical, Communication, and Computer Engineering (ICEET) in 2023 and 2024, and as Subcommittee Chair for Quantum Science and Technology at PIERS 2025. Dr. Qasymeh is a frequent invited speaker at global scientific forums and actively contributes to international research and development initiatives.



Shabbir Hussain is a PhD scholar in the Department of Mathematics, University of Lahore. under the supervision of Dr. Romana Ashraf. His research areas include nonlinear partial differential equations, bifurcation theory, chaos analysis, soliton dynamics,

and analytical solution methods for complex nonlinear systems. He has more than 10 research papers in very good journals. He has given many talks as invited speaker in many national conferences.



Farrah Ashraf is an assistant professor at The University of Lahore, Lahore, Department of Mathematics and Statistics. Her research areas are Nonlinear Partial Differential Equations, Soliton Theory, Numerical Methods, Mathematical Modeling, Biomathematics

and Fractional Differential Equations. She has published 21 research papers in internationally reputed journals, which have been extensively cited by researchers both nationally and internationally, with over 326 citations. She has a strong research profile, reflected by an h-index of 8 and an i10-index of 7. She has received the Research Productivity Award 2023 by The University of Lahore, Lahore.