

Determination of Radon-222 Levels, Concentration Level, and Its Health Hazards in Drinking Water from Gashua and Damaturu, Yobe State, Nigeria

Iliya I. Fatima*, Umaru Ibrahim, Inusa I. Ewa, Idris M. Mustapha, and Adamu Hussaini

¹Department of Physics, Nasarawa State University, Keffi, Nigeria.

¹Radiation and Medical Physics Unit, Department of Physics, Nasarawa State University, Keffi, Nigeria.

²Nigerian Nuclear Regulatory Authority, Abuja, Nigeria.

Received: 30 Aug. 2025, Revised: 20 Oct. 2025, Accepted: 30 Nov. 2025

Published online: 1 Jan. 2026

Abstract: Water quality is a vital component of public health and environmental safety, as clean water is essential for drinking, food production, and sanitation. In this study, the health risk assessment of heavy metals and Radon-222 levels in portable drinking water from Gashua and Damaturu, Yobe State, Nigeria, was investigated. Forty (40) samples (twenty (20) each from Damaturu and Gashua, respectively) were collected by simple random sampling and were analyzed for radon-222 concentration level Liquid Scintillation Spectrometer. Each of the samples was coded with a number for sample identification, and the GPS location of each sample was taken. The calculated Radon-222 concentration, annual effective dose by ingestion, dose contribution to the stomach due to ingestion, annual effective dose by inhalation estimation, dose contribution to the lung due to inhalation, and excess lifetime cancer risk for Damaturu (Gashua) are 780.61655 Bq/L (753.2699 Bq/L), 0.00569855 mSv/y (0.005499 mSv/y), 0.0006816 mSv/y (0.000658 mSv/y), 34.34713 mSv/y (33.14388 mSv/y), 4.1182207 mSv/y (3.973951 mSv/y), and 1.99448E-05 (1.92E-05), respectively. The result of radon-222 concentration indicates an elevated radon concentration in water, which poses potential radiological health risks through both ingestion and inhalation exposure. However, the computed excess lifetime cancer risk (1.92E-05) falls within the acceptable risk range of 10^{-6} to 10^{-3} recommended by USEPA. Adequate portable drinking water monitoring is recommended, and safety measures should be put in place.

Keywords: Radon-222 concentration, liquid scintillation detector, radiological hazards, and well water.

1 Introduction

Radon-222 (^{222}Rn) is a radioactive noble gas that forms naturally during the decay process of uranium-238, commonly found in various geological formations [1- 5]. As an inert, invisible, and odorless gas, its presence in environmental media often goes unnoticed without analytical investigation. Although most studies on radon exposure focus on inhalation as the primary route—particularly from indoor air accumulation—emerging evidence highlights that radon dissolved in groundwater also constitutes a significant public health concern, especially in regions that depend heavily on such water sources for daily use [6-10].

In Yobe State, northeastern Nigeria, the towns of Gashua and Damaturu rely primarily on groundwater accessed through wells and boreholes. The underlying geology in this area may contain uranium-bearing rocks that release

radon into aquifers, thereby elevating its concentration in drinking water. When consumed over time, this radioactive element can accumulate in internal organs, where its decay products emit alpha particles that may damage cells and increase the risk of cancers, particularly in the stomach and gastrointestinal tract [11-15].

A study by Obed et al. [5] measured radon concentrations in groundwater across parts of southwestern Nigeria and found levels exceeding the recommended safety limits set by international agencies. The authors emphasized the need for routine monitoring and public awareness regarding radiation exposure through drinking water. Similarly, Arogunjo et al. [4] assessed the radiological quality of water sources in Nigeria and identified radon-222 as a notable contributor to the overall radiation dose received by the population. Their research underlined the significance of geological formations in influencing radionuclide

*Corresponding author e-mail: binafaibrahim@yahoo.com

content in water and recommended further assessments across other geopolitical zones.

Given the limited data on radon contamination in the northeastern region, this study seeks to quantify the concentration of radon-222 in drinking water sources in Gashua and Damaturu. In addition, it evaluates the potential health implications based on globally accepted radiological health benchmarks. The outcomes of this research are expected to support evidence-based environmental management policies and promote public health safety in the affected communities.

2 Materials and Methods

Study Area

Damaturu and Gashua are key towns in Yobe State, northeastern Nigeria, both situated in a semi-arid climate with hot, dry seasons and limited rainfall. Damaturu, located at approximately 11.7461° N, 11.9608° E, serves as the state capital, while Gashua, located at 12.8734° N, 11.0406° E, lies further north near the Komadugu Yobe River. Damaturu has a population of around 44,000, whereas Gashua has about 125,000 residents.

Agriculture is central to the economy in both towns, with residents engaged in cultivating crops such as millet, sorghum, maize, and rice. Gashua, benefiting from the nearby river, supports irrigation farming, which allows for year-round crop production. In addition to farming, livestock rearing, including cattle, sheep, and goats, is common, especially among the Kanuri and Fulani ethnic groups.

Both towns face challenges, including environmental issues like desertification and climate impacts on water resources. Security issues have also disrupted economic stability, but recent improvements have enabled slow recovery efforts. The people's resilience continues to drive progress in these communities, even amid ongoing challenges.




Fig. 1: Map of Study Area.

Population, Sample, and Sampling Technique

The population of the study covers drinking water sources from Gashua and Damaturu, Yobe State, Nigeria. For the purpose of obtaining representative water samples and to ensure that all water samples have equal chances of selection, a simple random sampling technique was adopted to collect forty (40) water samples from the study area.

Method of Sample Collection

The plastic containers were first washed and rinsed with distilled water to avoid radon present in the samples from being contaminated. Water samples were preserved with 20 ml of concentrated HNO₃ per liter of the water in order to minimize absorption of the dissolved radon on container walls.

The water samples were collected from the well to ensure fresh samples were obtained. At each location, the containers were filled to the brim with the water sample without any headspace to prevent CO₂ from being trapped and dissolving in water, which might affect the chemistry, e.g., pH, and then immediately closed to avoid loss of radon by degassing during transport to the laboratory.

The samples were sent for analysis immediately after collection without allowing them to stay long (three days maximum) to minimize the influence of radioactive decay. This will be done to achieve maximum accuracy and not to alter its composition. Each of the samples will be coded with a number for simple identification. The sample codes with the GPS location of each sample were taken.

Method of Sample Preparation for Radon Concentration

10 ml each of the water samples was transferred into a 20 ml glass scintillation vial, to which 10 ml of Insta-Gel scintillation cocktail was added. It was then sealed tightly. The vials were shaken for more than two minutes to extract radon-222 in the water phase into the organic scintillate, and the samples collected were counted for 60 minutes in a liquid scintillation counter using energy discrimination for alpha particles.

Method of Sample Analysis for Radon-222

The glass scintillation vial will be sealed tightly. The vials were shaken for more than two minutes to extract radon-222 in the water phase into the organic scintillate, and the samples collected were counted for 60 minutes in a liquid scintillation counter using energy discrimination for alpha particles.

The samples were analyzed using a Liquid Scintillation Counter (Tri-Carb LSA 1000TR) model located at the Centre for Energy Research and Training (CERT), Ahmadu Bello University, Zaria-Nigeria.

Radon-222 Concentration in Bq/l Estimation Method

The activity concentration of Radon-222 was calculated from the samples and background results obtained using the formula below:

$$C_{Rn-222}(BqL^{-1}) = \frac{C-C_b}{CF \times DE} \cdot e^{\lambda t} \quad 1$$

where,

C_{Rn-222} is the Radon-222 activity concentration in water (BqL^{-1}), C is the measured count rate (cps) from the scintillation counter, C_b is the background count rate (cps), CF is the Calibration factor (0.2 cps per Bq/L), DE is the efficiency of the detector (0.08), t is the Elapsed time (in days) between sample collection and measurement, and λ is the decay constant of radon-222 ($1.26 \times 10^{-4} \text{ min}^{-1}$).

3.2.7.2 Annual Effective Dose by Ingestion Estimation Method

The corresponding annual effective doses (mSv/y) due to ingestion of Radon-222 in water samples from the study area were also calculated using equation 3.2 by taking into account the dose coefficient in (Sv/Bq), the annual water consumption (L/Y), and the activity concentration of Radon-222 obtained from equation 3.1 using equation 2 by Rilwan et al. [16];

$$E = C_{Rnw} \times D \times L \quad 2$$

where: C_{Rnw} is the concentration of Radon-222, D is the dose coefficient ($10^{-8} Sv/Bq, 2 \times 10^{-8} Sv/Bq, 7 \times 10^{-8} Sv/Bq$) for adults, children, and infants, respectively. L is the annual water consumption by an adult of 2 litres per day, that is $730L/Y$ [17].

According to the United Nations Scientific Committee on the Effects of Atomic Radiation [17], doses due to ingestion of radon in water for similar consumption rates could be a factor of 2 and 7 higher for children and infants, respectively.

3.2.7.3 Dose Contribution to the Stomach due to Ingestion

It is a product of the stomach tissue weighted factor (0.1196) with the corresponding ingestion dose using Equation (3) below

$$D_{stomach} = E_{ing} \times 0.1196 \quad 3$$

3.2.7.4 Annual Effective Dose by Inhalation Estimation Method

The annual effective dose of Radon by inhalation was estimated using equation (4) as:

$$He = C \times F \times T \times D \quad 4$$

He is the annual effective dose (mSv^{-1}), C is the radon concentration (BqL^{-1}), F is the Equilibrium factor (0.5), T is the Indoor occupancy time (8000 h^{-1}), and D is the dose conversion factor ($1.1 \times 10^{-5} \text{ mSvh}^{-1}/BqL^{-1}$) [18-21].

3.2.7.5 Dose Contribution to the Lung due to Inhalation

It is the product of the lung tissue weighted factor (0.1199) and the corresponding inhalation dose using Equation (5)

$$D_{lung} = He \times 0.1199 \quad 5$$

3.2.7.6 Excess Lifetime Cancer Risk Estimation Method

The excess lifetime cancer risk was evaluated using equation (6) as:

$$ELCR = AEDE \times DL \times RF \times 10^{-3} \quad 6$$

where $ELCR$ is the excess lifetime cancer risk, $AEDE$ is the annual effective dose equivalent, DL is the average duration of life (estimated to 70 years), and RF is the Risk Factor (Sv^{-1}), i.e., fatal cancer risk per Sievert. For stochastic effects, ICRP uses RF as 0.05 for the public [1].

3 Results and Discussion

The results of radon-222 concentration in water samples (in CPMC) from Gashua and Damaturu, Yobe State, Nigeria, are presented in Figures 2 and 3, respectively.




Fig.1: Radon-222 concentration in count per minute of water sample collected from Gashua, Yobe State, Nigeria.




Fig.2: Rado-222 concentration in count per minute of water sample collected from Damaturu, Yobe State, Nigeria.

Figures 1 and 2 present the raw results of Rado-222 concentration in count per minute of water samples collected from Gashua and Damaturu, respectively. The raw result of Radon-222 concentration in counts per minute (cpm) represents the detected radioactivity from a sample, as measured by a radiation detector such as a scintillation counter, Geiger-Müller tube, or an alpha spectrometer. The cpm value indicates the number of radiation events (alpha decays) detected per minute and depends on factors like detector efficiency, background radiation, and measurement conditions.

The calculated hazard parameter due to radon-222 concentration level in water, and the background results obtained were determined. The annual effective dose by ingestion, dose contribution to the stomach due to ingestion, annual effective dose by inhalation estimation, dose contribution to the lung due to inhalation, and excess lifetime cancer risk, were estimated. Table 1 and 2 present the result of the calculated hazard parameters of radon-222 concentration in water sample collected from Damaturu, and Gashua, respectively.

Table 1: Calculated hazard parameters of radon-222 concentration in water sample collected from Damaturu, Yobe State.

Sample Code	C_{Rn-222} (Bq/L)	E_{Ing} (mSv/y)	$D_{Stomach}$ (mSv/y)	E_{inh} (mSv/y)	D_{Lung} (mSv/y)	ELCR
DAM1	1128.949	0.008241	0.000986	49.67378	5.955886	2.88447E-05
DAM2	999.2644	0.007295	0.000872	43.96763	5.271719	2.55312E-05
DAM3	1412.753	0.010313	0.001233	62.16114	7.45312	3.60958E-05
DAM4	340.8149	0.002488	0.000298	14.99586	1.798003	8.70782E-06
DAM5	1023.698	0.007473	0.000894	45.0427	5.40062	2.61555E-05
DAM6	836.3749	0.006106	0.00073	36.8005	4.412379	2.13694E-05
DAM7	380.9108	0.002781	0.000333	16.76008	2.009533	9.73227E-06
DAM8	592.0407	0.004322	0.000517	26.04979	3.12337	1.51266E-05
DAM9	432.9101	0.00316	0.000378	19.04805	2.283861	1.10609E-05
DAM10	1130.829	0.008255	0.000987	49.75647	5.965801	2.88927E-05
DAM11	586.4022	0.004281	0.000512	25.8017	3.093623	1.49826E-05

DAM12	786.8815	0.005744	0.000687	34.62279	4.151272	2.01048E-05
DAM13	568.8602	0.004153	0.000497	25.02985	3.001079	1.45344E-05
DAM14	848.9048	0.006197	0.000741	37.35181	4.478482	2.16895E-05
DAM15	1033.722	0.007546	0.000903	45.48376	5.453503	2.64116E-05
DAM16	984.2284	0.007185	0.000859	43.30605	5.192395	2.5147E-05
DAM17	507.4634	0.003704	0.000443	22.32839	2.677174	1.29657E-05
DAM18	882.1092	0.006439	0.00077	38.81281	4.653656	2.25379E-05
DAM19	354.5979	0.002589	0.00031	15.60231	1.870717	9.05998E-06
DAM20	780.6166	0.005699	0.000682	34.34713	4.118221	1.99448E-05
Mean	780.61655	0.00569855	0.0006816	34.34713	4.1182207	1.99448E-05

Table 2: Calculated hazard parameters of radon-222 concentration in water sample collected from Gashua, Yobe State.

Sample Code	C _{Rn-222} (Bq/L)	E _{Ing} (mSv/y)	D _{Stomach} (mSv/y)	E _{inh} (mSv/y)	D _{Lung} (mSv/y)	ELCR
GSH1	403.4647393	0.002945293	0.000352257	17.75244853	2.128518579	1.03085E-05
GSH2	1973.468834	0.014406322	0.001722996	86.83262868	10.41123218	5.04221E-05
GSH3	479.8975005	0.003503252	0.000418989	21.11549002	2.531747254	1.22614E-05
GSH4	903.4101772	0.006594894	0.000788749	39.7500478	4.766030731	2.30821E-05
GSH5	342.0679312	0.002497096	0.000298653	15.05098897	1.804613578	8.73984E-06
GSH6	395.3202648	0.002885838	0.000345146	17.39409165	2.085551589	1.01004E-05
GSH7	1109.528033	0.008099555	0.000968707	48.81923346	5.853426092	2.83484E-05
GSH8	559.4627519	0.004084078	0.000488456	24.61636108	2.951501694	1.42943E-05
GSH9	946.0120441	0.006905888	0.000825944	41.62452994	4.99078114	2.41706E-05
GSH10	1068.179162	0.007797708	0.000932606	46.99988314	5.635285989	2.7292E-05
GSH11	711.0752782	0.00519085	0.000620826	31.28731224	3.751348738	1.8168E-05
GSH12	380.9108098	0.002780649	0.000332566	16.76007563	2.009533068	9.73227E-06
GSH13	466.7410416	0.00340721	0.000407502	20.53660583	2.462339039	1.19252E-05
GSH14	432.9101473	0.003160244	0.000377965	19.04804648	2.283860773	1.10609E-05
GSH15	1068.179162	0.007797708	0.000932606	46.99988314	5.635285989	2.7292E-05
GSH16	586.4021677	0.004280736	0.000511976	25.80169538	3.093623276	1.49826E-05
GSH17	598.9321286	0.004372205	0.000522916	26.35301366	3.159726338	1.53027E-05
GSH18	568.8602225	0.00415268	0.00049666	25.02984979	3.00107899	1.45344E-05
GSH19	974.204456	0.007111693	0.000850558	42.86499606	5.139513028	2.48909E-05
GSH20	1096.371574	0.008003512	0.00095722	48.24034927	5.784017877	2.80123E-05
Mean	753.2699	0.005499	0.000658	33.14388	3.973951	1.92E-05

The annual effective dose due to ingestion of Damaturu (Gashua) is 0.00569855 mSv/y (0.005499 mSv/y), which is below the International Commission on Radiological Protection (ICRP) reference dose limit of 0.1 mSv/y for drinking water [27-29]. This indicates that the ingestion route does not contribute significantly to radiation exposure. Similarly, the stomach dose contribution (0.0006816 mSv/y) is minimal, suggesting that ingestion of radon-contaminated water is not the primary concern.

The estimated annual effective dose by inhalation of Damaturu (Gashua) is 34.34713 mSv/y (33.14388 mSv/y) is alarmingly high compared to the ICRP recommended public exposure limit of 1 mSv/y [13]. This level is of significant concern because inhalation is the primary route of radon exposure, leading to an increased risk of lung cancer. The lung dose contribution (4.1182207 mSv/y) further supports the potential radiological impact on respiratory health.

The computed excess lifetime cancer risk of Damaturu (Gashua) is 1.99448E-05 (1.92E-05) is within the acceptable range of 10^{-6} to 10^{-3} recommended by USEPA [22]. However, considering the high inhalation dose, there is a need for risk mitigation strategies to reduce radon exposure.

4 Conclusion

The study provides results of radon-222 contamination in some selected well water sources from Gashua and Damaturu, Yobe State, Nigeria. Radon-222 concentrations in groundwater were found to be significantly high, surpassing international safety guidelines. The major risk is through inhalation, leading to a high annual effective dose and increased lung cancer risk. The estimated excess lifetime cancer risk (ELCR) due to radon exposure is above acceptable limits, suggesting an increased probability of developing lung cancer over a lifetime for individuals who consume and use the contaminated water regularly. Therefore, proper mitigation strategies should be carried out so that communities in Gashua and Damaturu may not face increasing health burdens related to chronic heavy metal toxicity and radiation exposure.

References

- [1] Ingawa A. F., Umaru I., Mundi A. A., Idris M. M., Isa S., and Ingawa A. F. (2025). Radiometric Evaluation of Radon-222 Concentration in Portable Drinking Water Around Proposed Crude Oil Drilling Site in Obi, Nasarawa State, Nigeria. *Dutse Journal of Pure and Applied Sciences (DUJOPAS)*, 11(2), 116-132.
- [2] Abdullahi A. M., Idris M. M., & Rabo M. (2021). Assessment of Outdoor Gamma Exposure Level of Some Swampy Agricultural Soils of Nasarawa West, Nigeria. *Asian Journal of Research and Reviews in Physics*, 5(2), 32-41. <https://www.journalajr2p.com/index.php/AJR2P/article/view/30160/56585>
- [3] Abubakar, S. B., Usman, R., Ibrahim, U., Samson, D. Y., Idris, M. M., Abdullahi, A. M., & Ibrahim, M. (2023). Toxicity of Radon-222 in Groundwater across Keana in Nasarawa, Nigeria. *Advances in Geological and Geotechnical Engineering Research*, 5(2), 38- 49. <https://journals.bilpubgroup.com/index.php/agger>
- [4] Arogunjo, A. M., Farai, I. P., & Fuwape, I. A. (2005). Contribution of natural radionuclides to the terrestrial gamma dose rate in southwestern Nigeria. *Radiation Protection Dosimetry*, 114(4), 530–532. <https://doi.org/10.1093/rpd/nci282>
- [5] Obed, R. I., Ademola, J. A., & Popoola, O. E. (2012). Radon concentration in groundwater in selected areas of southwestern Nigeria and the associated health risk. *Radiation Protection Dosimetry*, 150(4), 515–519. <https://doi.org/10.1093/rpd/ncr404>
- [6] World Health Organization (WHO). (2011). *Guidelines for drinking-water quality* (4th ed.). WHO Press.
- [7] Akinyemi, O., Olowofela, J. A., & Akinpelu, A. (2020). Environmental implications of naturally occurring radioactive materials in oil-producing regions. *Journal of Environmental Radioactivity*, 215, 106176. <https://doi.org/10.1016/j.jenvrad.2020.106176>
- [8] International Atomic Energy Agency. (2018). *Guidelines on radiological monitoring of drinking water sources*. IAEA Safety Report Series.
- [9] Atef El-Taher, HMH Zakaly., 2020 Radon concentration and radon exhalation rate for basement rocks from Wadi Um Huytat in Central Eastern Desert of Egypt. *AIP Conference Proceedings* 2313 (1), 020006.
- [10] F Alshahri, Atef El-Taher, AEA Elzain., 2019 Measurement of radon exhalation rate and annual effective dose from marine sediments, Ras Tanura, Saudi Arabia, using CR-39 detectors. *Romanian Journal of Physics*. 64, 811.
- [11] Mostafa Y.A. Mostafa and Atef El-Taher., 2019 Radon Standard Source in Different Countries with Different Principals. *J. Rad. Nucl. Appl.* 4, No. 1, 35-41.
- [12] Fatimh Alshahri, Atef El-Taher, and Abd Elmoniem Ahmed Elzain., 2019 Measurements of radon exhalation rate and annual effective dose from Marine sediments, Ras Tannura, Saudi Arabia, using CR-39 detectors. *Romanian Journal of Physics* 64-811:1-13
- [13] Fatimh Alshahri, Atef El-Taher, and Abd Elmoniem Ahmed Elzain., 2017 Characterization of Radon

- Concentration and Annual Effective Dose of Soil Surrounding a Refinery Area, Ras Tanura, Saudi Arabia. *Journal of Environmental Science and Technology*, 10(6):311-319
- [14] Manar Dheyaa Salim, Ali A. Ridha, Nada Farhan Kadhim, and Atef El- Taher., 2020 Effects of Changing the Exposure Time of CR-39 Detector to Alpha Particles on Etching Conditions. *J. Rad. Nucl. Appl.* 5, No. 2, 119-125.
- [15] A. El-Taher, M El-Hagary, M Emam-Ismail, FA El-Saied, FA Elgendy, Radon and its decay products in the main campus of Qassim University, Saudi Arabia, and its radiation hazards. *Journal of American Science* 9 (6), 257-266. 2013.
- [16] Rilwan, U., Yahaya, I., Musa, M., Galadima, O.O., Waida, J., & Idris, M. M. (2022). Investigation of Radon-222 in Water from Loko Town in Nasarawa, Nasarawa State, Nigeria. *Journal of Oncology and Cancer Screening*, 4(2), 1-7. DOI: <http://doi.org/005.2022/1.1053>. <https://aditum.org/journals/oncology-and-cancer-screening>
- [17] United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) (2000). *Sources and Effects of Ionizing Radiation*. Report to General Assembly, (Vol. 1). Annex B, United Nations, New York.
- [18] A. El-Taher and Ahmed El-Turki., 2016 Radon Activity Measurements in Irrigation Water from Qassim Province Using RAD7. *Journal of Environmental Biology* 37, 1299-1302.
- [19] Atef El-Taher, LA Najam, AA Abojassim, H Mraity., 2020 Assessment of Annual Effective Dose for Different Age Groups Due to Radon Concentrations in Groundwater Samples at Qassim, Saudi Arabia. *Iranian Journal of Medical Physics* 17: 15-2
- [20] Wedad Rayif Alharbi, Adel G. E. Abbady ,and A. El-Taher., 2014 Radon Concentrations Measurement for groundwater Using Active Detecting Method American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS) 14 (1) 1-11.
- [21] S. S. Althoyaib and A. El-Taher., 2014 The Measurement of Radon and Radium Concentrations in well water from Al-Jawa, Saudi Arabia, *Journal of Natural Sciences and Mathematics Qassim University*, Vol. 7, No. 2, PP 179-192.
- [22] United States Environmental Protection Agency (USEPA). (1991). National primary drinking water regulations for radionuclides. *Federal Register*, 56(138), 33050-33127.
- [23] A. El-Taher., 2012 Measurement of Radon Concentrations and Their Annual Effective Dose Exposure in Groundwater from Qassim Area, Saudi Arabia. *Journal of Environmental Science and Technology* 5 (6) 475-481.
- [24] E. E. Massoud, Atef El-Taher, Abd Elmoniem A. Elzain., 2020 Estimation of environmental radioactivity and Radiation Dose from Exposure to Radon in groundwater for inhabitants in Qassim area - Saudi Arabia. *Desalination and Water Treatment*, 205, 308–315.
- [25] Atef El-Taher., 2018 An Overview of Instrumentation for Measuring Radon in Environmental Studies. *J. Rad. Nucl. Appl.* 3, No. 3, 135-141.
- [26] K Said, SM Brahim, EB Abderrahman, A. El-Taher., 2024 Determination of Radon Gas Concentration in the Water of Midelt Region, Morocco, Using a Nuclear Track Detector (LR-115) and Assessment of Radiological Health Risk. *Pollution* 10 (1), 248-255.
- [27] ICRP (2007). *Annals of the International Commission on Radiological Protection*. Publication, 23.
- [28] AS Bako, I Umar, SD Yusuf, U Rilwan, MM Idris, NM Abdu, AM El-Taher., 2023 Determination of Effective Dose Due to Radon Ingestion and Inhalation from Groundwater Across Awe Local Government Areas in Nasarawa, Nigeria. *Rafidain Journal of Science* 32 (4), 92-105.
- [29] Khadour Said, El-Bokili Abderrahman, Atef El-Taher., 2023, Characterization of radon concentration, surface and mass exhalation rates radon and effective dose rate from soil samples in the Midelt, region of Morocco, using Alpha GUARD. *Journal of Wuhan University of Science and Technology*, 47(1):899-907.